arXiv:1503.04156v4 [math.DS] 26 Dec 2016

LOCAL LIMIT THEOREM IN NEGATIVE CURVATURE

FRANÇOIS LEDRAPPIER AND SEONHEE LIM

Abstract. Consider the heat kernel $\wp(t,x,y)$ on the universal cover $\widetilde{M}$ of a closed Riemannian manifold of negative curvature. We show the local limit theorem for $\wp$:

$$\lim_{t\to\infty} t^{3/2}e^{\lambda_0 t}\wp(t,x,y)=C(x,y),$$

where λ_0 is the bottom of the spectrum of the geometric Laplacian and $C(x,y)$ is a positive $(-\lambda_0)$-eigenfunction which depends on $x,y\in\widetilde{M}$.

We show that the λ_0-Martin boundary of $\widetilde{M}$ is equal to its topological boundary. The Martin decomposition of $C(x,y)$ gives a family of measures $\{\mu_x^{\lambda_0}\}$ on $\partial\widetilde{M}$. We show that $\{\mu_x^{\lambda_0}\}$ is the unique family minimizing the energy or the Rayleigh quotient of Mohsen.

We use the uniform Harnack inequality on the boundary $\partial\widetilde{M}$ and the uniform three-mixing of the geodesic flow on the unit tangent bundle SM for suitable Gibbs-Margulis measures.

1. Introduction

Let (M,g) be an m-dimensional closed connected Riemannian manifold of negative curvature, and $(\widetilde{M},\widetilde{g})$ its universal cover endowed with the lifted Riemannian metric. Let us denote by d the distance on M, $\widetilde{M}$, as well as on their unit tangent bundles $\pi : SM\to M$ and $\pi : S\widetilde{M}\to\widetilde{M}$ (see [**PPS**] for various distances on M and on SM and the equivalences between them). We denote the natural projection $(\widetilde{M},\widetilde{g})\to(M,g)$ and its derivative by p. The fundamental group $\Gamma=\pi_1(M)$ acts on $\widetilde{M}$ as isometries such that $M=\widetilde{M}/\Gamma$. Let M_0 be a bounded fundamental domain for this action.

We consider the Laplace-Beltrami operator $\Delta := \mathrm{Div}\nabla$ for smooth functions on $\widetilde{M}$ and the corresponding heat kernel function $\wp(t,x,y), t\in\mathbb{R}_+, x,y\in\widetilde{M}$, which is the probability density defined as the fundamental solution of the heat equation, i.e. the function which satisfies $\frac{\partial\wp}{\partial t}=\Delta_y\wp$ and $\lim\limits_{t\to 0}\wp(t,x,y)=\delta(x-y)$. The function $\wp$ is clearly Γ-invariant and symmetric.

Denote by λ_0 the bottom of the spectrum of the geometric Laplacian $-\Delta$ on $L^2(\widetilde{M},\mathrm{Vol})$, where $d\mathrm{Vol}(z)$ is the Riemannian volume form on $\widetilde{M}$. Since Γ is not amenable, λ_0 is

---

Date: April 6, 2016.

2000 *Mathematics Subject Classification.* Primary 37D40; 37A17; 37A25; 37A30; 37A50 .

Key words and phrases. local limit theorem, Brownian motion, rate of mixing.

positive [**Br**]. For all $x, y \in \widetilde{M}$, we have

$$\lambda_0 \;=\; \lim_{t\to\infty} -\frac{1}{t}\ln \wp(t,x,y) \tag{1.1}$$

by the spectral theorem.

Our main result is a local limit theorem which refines (1.1).

Theorem 1.1 (Local Limit Theorem)**.** *There exists a positive function C on $\widetilde{M}\times\widetilde{M}$ such that for all $x, y \in M$,*

$$\lim_{t\to\infty} t^{3/2} e^{\lambda_0 t}\wp(t,x,y) \;=\; C(x,y). \tag{1.2}$$

When $\widetilde{M}$ is the hyperbolic space $\mathbb{H}^3$, there is an explicit expression for $\wp(t,x,y)$ ([**DGM**]) and Theorem 1.1 is clear, with

$$C(x,y) = (4\pi)^{-3/2}\frac{d(x,y)}{\sinh d(x,y)}.$$

More generally, if $\widetilde{M}$ is a symmetric space of non-compact type, then Theorem 1.1 with $t^{k/2}$ instead of $t^{3/2}$ was proven by Bougerol ([**B**]) using representation theory of the group G of orientation-preserving isometries of $\widetilde{M}$. The integer k is given by the rank plus twice the number of positive indivisible roots, which depends only on the Lie algebra of the semi-simple group G. In particular, $k = 3$ for all rank one symmetric spaces and this explains why one might expect $t^{3/2}$ for negatively curved manifolds. Bougerol proved the theorem for all random walks on semi-simple groups with a distribution that is left and right K-invariant, which implies the same result for Brownian motions on $\widetilde{M} = G/K$.

The limit function $C(x,y)$ is symmetric by Theorem 1.1 and it is a positive $(-\lambda_0)$-eigenfunction in y: we further give a formula in Theorem 1.7 below. However, it was already known that if the limit

$$\lim_{t\to\infty}\frac{\wp(t,x,y)}{\wp(t,x,x)} \;=\; \frac{C(x,y)}{C(x,x)} \tag{1.3}$$

exists on a Riemannian manifold, then $C(x,y)$ is a $(-\lambda_0)$-eigenfunction in y [**ABJ**] (Theorem 1.2). It is indeed a conjecture by Davies ([**Da**]) that the limit (1.3) always exists (see [**Ko**] for a recent counter-example for the analogous question on graphs). Our result can be stated as:

Corollary 1.2. *The universal cover of a compact Riemannian manifold with negative sectional curvature satisfies Davies conjecture.*

See [**ABJ**] for further discussion and applications of Davies conjecture.

A similar local limit theorem was first observed by Gerl [**Ge**] and Woess [**GW**] for random walks on a free group which are supported by the generators of the group. It was then proven by Lalley for random walks with finite support on a finitely generated free group [**La**]. This was extended by Gouëzel and Lalley to symmetric random walks with finite support on cocompact Fuchsian groups [**GL**] and finally by Gouëzel to symmetric random walks with finite support on hyperbolic groups [**G1**]. Our proof follows the strategy and ideas of [**GL**] and [**G1**]. By [**G2**], this general strategy works for measures of infinite support and with superexponential moments.

Two main new ingredients of the proof of Theorem 1.1 are the uniform rapid-mixing of the geodesic flow generalizing Dolgopyat theorem and the generalised Patterson-Sullivan conformal family whose Radon-Nikodym derivative is the Martin kernel $k^2_{\lambda_0}(x,y,\xi)$, which is defined in Theorem 1.4 below.

As in [**G1**], we obtain several subsequent results which have their own interest. Let us introduce more notation to describe these results. For any real $\lambda < \lambda_0$, we define the λ-*Green function* G_λ: for all $x \neq y \in \widetilde{M}$,

$$G_\lambda(x,y) \ := \ \int_0^\infty e^{\lambda t} \wp(t,x,y)dt.$$

The integral converges at ∞ thanks to the spectral theorem (1.1). The integral converges at 0 since as $t \to 0$, $\wp(t,x,y) \sim C/t^{m/2} e^{-\frac{d^2(x,y)}{4t}}$. The function $G_\lambda(x,\cdot)$ is positive $(\Delta + \lambda)$-harmonic outside $\{x\}$.

Guivarc'h observed that the λ_0-Green function is finite as well: for all $x \neq y \in \widetilde{M}$,

$$G_{\lambda_0}(x,y) \ := \int_0^\infty e^{\lambda_0 t} \wp(t,x,y)dt < \infty.$$

Moreover, for any compact $K \subset \widetilde{M}$, we also have

$$\int_K G_{\lambda_0}(x,y)\, d\mathrm{Vol}(y) \ < \ \infty. \tag{1.4}$$

(See Lemma 2.1.)

In Section 3, we show (see Proposition 3.14, where we relate τ with other dynamical properties)

Theorem 1.3. *There are positive constants τ and C such that, for $x, y \in \widetilde{M}$ with $d(x,y) \geq 1$,*

$$G_{\lambda_0}(x,y) \ \leq \ Ce^{-\tau d(x,y)}.$$

Two geodesic rays in $\widetilde{M}$ are said to be *equivalent* if they remain a bounded distance apart. The *geometric boundary* $\partial\widetilde{M}$ is defined as the space of equivalence classes of unit speed geodesic rays. A sequence $\{y_n\}_{n\in\mathbb{N}}$ in $\widetilde{M}$ converges to a point in $\partial\widetilde{M}$ if, and only if, for some (hence, for all) $x \in \widetilde{M}$,

$$d(x,y_n) + d(x,y_m) - d(y_n,y_m) \to \infty \quad \text{as} \ \ n,m \to \infty.$$

We now describe the Martin boundary of the operator $\Delta + \lambda_0$.

Theorem 1.4. *[λ_0-Martin boundary] Fix $x \in \widetilde{M}$ and assume that the sequence $\{y_n\}_{n\in\mathbb{N}}$ converges to a point $\xi \in \partial\widetilde{M}$. Then, there exist a positive $(-\lambda_0)$-eigenfunction $k_{\lambda_0}(x,y,\xi)$ of the Laplacian, which we call Martin kernel, such that*

$$\lim_{n\to\infty} \frac{G_{\lambda_0}(y,y_n)}{G_{\lambda_0}(x,y_n)} \ = \ k_{\lambda_0}(x,y,\xi).$$

Moreover, the Martin boundary of $\Delta + \lambda_0$ *coincides with the geometric boundary. In particular, for any positive harmonic function* F *of* $\Delta + \lambda_0$*, there is a finite measure* ν_F *on* $\partial \widetilde{M}$*, depending on* x*, such that*

$$F(y) \ = \int_{\partial \widetilde{M}} k_{\lambda_0}(x, y, \xi) d\nu_F(\xi).$$

See Section 4 for the proof and more properties of the Martin kernel $k_{\lambda_0}(x, y, \xi)$. Martin kernel plays the role of a conformal density for a family of measures on the boundary $\partial \widetilde{M}$.

Theorem 1.5. *For all* $y \in \widetilde{M}$*, there is a finite measure* $\mu_y^{\lambda_0}$ *on* $\partial \widetilde{M}$ *such that*

1) *the family* $y \mapsto \mu_y^{\lambda_0}$ *is* Γ*-equivariant:* $\mu_{\gamma y}^{\lambda_0} = \gamma_*(\mu_y^{\lambda_0})$ *for* $\gamma \in \Gamma$ *and*
2) *for* $\mu_x^{\lambda_0}$*-a.e.* $\xi \in \partial \widetilde{M}$*, all* $y \in \widetilde{M}$*,*

$$\frac{d\mu_y^{\lambda_0}}{d\mu_x^{\lambda_0}}(\xi) \ = \ k^2_{\lambda_0}(x, y, \xi).$$

The family is unique if we normalize by $\int_M \mu_x^{\lambda_0}(\partial \widetilde{M}) d\,Vol(x) = 1$.

For any Γ-equivariant families $\nu = \{\nu_x\}_{x \in \widetilde{M}}$ of measures on $\partial \widetilde{M}$ with cocycle $k(x, y, \xi) := \frac{d\nu_y}{d\nu_x}(\xi)$ and normalized by $\int_{M_0} \nu_x^{\lambda_0}(\partial \widetilde{M}) d\mathrm{Vol}(x) = 1$, we define the *energy of* ν as follows:

$$\mathcal{E}(\nu) := \int_{M_0} \left(\int_{\partial \widetilde{M}} \|\nabla_{y=x} \ln k(x, y, \xi)\|^2 d\nu_x(\xi) \right) d\mathrm{Vol}(x),$$

We define the energy to be infinite if the gradient is not defined almost everywhere.

The energy is equal to 4 times the Rayleigh quotient

$$\mathcal{R}(\nu) := \int_{M_0} \left(\int_{\partial \widetilde{M}} \|\nabla_x \sqrt{k(x_0, x, \xi)}\|^2 d\nu_{x_0}(\xi) \right) d\mathrm{Vol}(x)$$

defined by O. Mohsen in [**Mo**]. Mohsen showed that $\lambda_0 = \inf_\nu \mathcal{R}(\nu)$ and asked whether the minimum is achieved. Using Rayleigh quotient, it is easy to see that μ^{λ_0} has energy $\mathcal{E} = 4\lambda_0$.

Theorem 1.6. *The family* $\mu_x^{\lambda_0}$ *is the unique conformal family of measures which achieves the minimum energy.*

Mohsen proved the uniqueness for the manifolds with constant negative curvature.

The family $\mu_x^{\lambda_0}$ is a fourth natural Γ-equivariant family $\nu = \nu_x$ of measures on $\partial \widetilde{M}$ with regular cocycles, alongside with the Lebesgue visual measures, the Margulis-Patterson-Sullivan measures and the harmonic measures.[1] For rank one symmetric spaces, the Lebesgue measure on $\partial \widetilde{M}$ achieves the minimum. It is easy to see that the family $\mu_x^{\lambda_0}$ is proportional to the harmonic measures or to the Margulis-Patterson-Sullivan measures only when the manifold $\widetilde{M}$ is a symmetric space [**L3**].

---

[1] Observe that the energy of Margulis-Patterson-Sullivan measure is the volume entropy squared, and the energy of harmonic measure is Kaimanovich entropy [**H2**], [**K1**], [**L3**].

The last result we would like to emphasize is a formula of the function $C(x, y)$ in Theorem 1.1. In case of symmetric spaces, the function $C(x, y)$ is proportional to the positive $(-\lambda_0)$-eigenfunction invariant under the stabilizer K_x of the point x, a.k.a. the *Harish-Chandra function*, or the ground state, centered at x.

Theorem 1.7. *Fix $x \in \widetilde{M}$. There is a constant $\Upsilon = \Upsilon_{\lambda_0}$, given by (2.11), such that the positive $(-\lambda_0)$-eigenfunction $C(x, y)$ satisfies*

$$C(x,y) \;=\frac{\sqrt{\Upsilon}}{2\sqrt{\pi}}\int_{\partial\widetilde{M}} k_{\lambda_0}(x,y,\xi)d\mu_x^{\lambda_0}(\xi) \;=\; \frac{\sqrt{\Upsilon}}{2\sqrt{\pi}}\int_{\partial\widetilde{M}}\sqrt{d\mu_x^{\lambda_0}(\xi)}\sqrt{d\mu_y^{\lambda_0}(\xi)}.$$

Here, $\displaystyle\int_{\partial\widetilde{M}}\sqrt{d\mu_x^{\lambda_0}(\xi)}\sqrt{d\mu_y^{\lambda_0}(\xi)} := \int_{\partial\widetilde{M}}\sqrt{\frac{d\mu_y^{\lambda_0}}{d\mu_x^{\lambda_0}}(\xi)}d\mu_x^{\lambda_0}(\xi)$ as used in unitary representation of Γ associated to its action on $(\partial\widetilde{M}, \mu^{\lambda_0})$.

The article is organized along the path of the proof of Theorem 1.1.

In Section 2, we recall the consequences of Ancona's boundary Harnack inequality for $\lambda < \lambda_0$ ([**An1**]), in conjunction with the thermodynamic formalism for the geodesic flow (following [**K1**], [**H3**] and [**L2**]). Using mixing properties of the geodesic flow on the unit tangent bundle SM for suitable Γ-invariant Gibbs measures, we show that there is a function $P(\lambda)$ of λ and a positive function $D(x, \lambda)$ such that, for $\lambda < \lambda_0$, as $R \to \infty$

$$e^{-P(\lambda)R}\int_{S(x,R)} G_\lambda^2(x,z)dz \to D(x,\lambda), \tag{1.5}$$

where $S(x, R)$ is the sphere of radius R centered at x. (see Proposition 2.10).

We also recall from [**H3**] Corollary 5.5.1 that $\int_{S(x,R)} G_{\lambda_0}^2(x,z)dz$ is bounded independently from R (Proposition 2.12).

In Section 3, we use this bound to establish the uniform Harnack inequality at the boundary, i.e. the Ancona-Gouëzel inequality (Theorem 3.4). Theorem 1.4 follows and the other applications of thermodynamic formalism hold equally at $\lambda = \lambda_0$.

In Section 4, we discuss limits of measures on large spheres using uniform mixing of the geodesic flow. One consequence of our results is that the measures $\mu_{x,R}$ on the spheres $S(x, R)$ with density $e^{-RP(\lambda_0)}G_{\lambda_0}^2(x, y)$ converge to some measure $\mu_x^{\lambda_0}$ as $R \to \infty$ (Corollary 4.7). The measures $\mu_x^{\lambda_0}$ turns out to be a Γ-equivariant family with regular cocycle $e^{-P(\lambda_0)\beta(x,y,\xi)}k_{\lambda_0}^2(x, y, \xi)$, where $\beta(x, y, \xi)$ is the Busemann function (see the equation (2.7)). On the other hand, for $\lambda \in [0, \lambda_0], x \in \widetilde{M}$ and $R > 2$, we define the measure $m_{x,\lambda,R}$ on SM by first lifting the measure $e^{-P(\lambda)R}G_\lambda^2(x, z)dz$ on $S(x, R)$ to the set of unit vectors pointing towards x, and then projecting to SM by p. Another consequence is that there exists a probability measure $\overline{m}$ over SM such that the measures $m_{x,\lambda,R}$ converge towards some measure $\mu_x^{\lambda_0}(\partial\widetilde{M})\overline{m}$ on SM as $R \to \infty$ and $\lambda \to \lambda_0$ (see Corollary 4.8).

Once we prove that $P(\lambda_0)=0$ in Section 5, the family of measures $\mu_y^{\lambda_0}$ satisfies the statements of Theorem 1.5. We also obtain that for $x,y\in\widetilde{M}$, $\lim_{\lambda\to\lambda_0} -P(\lambda)\frac{\partial}{\partial\lambda}G_\lambda(x,y)$ is proportional to $C(x,y)$.

By a precise study of the second derivative $\frac{\partial^2}{\partial\lambda^2}G_\lambda(x,y)$ in Section 6.1, we obtain that both

$$\frac{P(\lambda)}{\sqrt{\lambda_0-\lambda}} \text{ and } \sqrt{\lambda_0-\lambda}\frac{\partial}{\partial\lambda}G_\lambda(x,y)$$

converge towards positive numbers as $\lambda\to\lambda_0$. In Section 6.2, we conclude the proof of Theorem 1.1 from Theorem 6.1 and a Tauberian Theorem as in [**GL**]. Theorem 1.7 follows as well.

In Section 7, we prove a uniform version of Dolgopyat's rapid-mixing for hyperbolic flows which is of independent interest. In Section 8, for completeness, we prove the precise balayage estimates in the form that is used in the article.

Remark 1.8. In this text, C stands for a number depending only on the geometry of M and Γ. However, its actual value may change from line to line. For the sake of clarity, we specify $C_0,\cdots,C_{11},C_\varepsilon,C(T)$ when the same number is used in another computation. Note that C_1,C_6,C_7 in Section 7 have the same role as in [**Me**]. Likewise, we consider spaces of α-Hölder continuous functions for some α of which the actual value may vary.

Acknowledgement : We would like to thank M. Pollicott for generously sharing his insights and ideas [**P1**], [**P2**], P. Bougerol for his interest and the [**ABJ**] reference and S. Gouëzel for helpful comments. The work was supported by University of Notre Dame, Seoul National University and MSRI during our visits. The second author was supported by NRF-2013R1A1A2011942, NRF-20100019516 and Korea Institute for Advanced Study (KIAS).

2. Potential theory and thermodynamic formalism

We recall in this section the results obtained by applying classical potential theory to the Laplacian on $\widetilde{M}$ and thermodynamic formalism to the geodesic flow.

Lemma 2.1. *For any compact set $K\subset\widetilde{M}$ with non-empty interior, we also have*

$$\int_K G_{\lambda_0}(x,y)\,d\mathrm{Vol}(y) \;<\; \infty. \tag{2.1}$$

Proof. Let ϕ be a $(-\lambda_0)$-eigenfunction of the Laplacian : $\Delta\phi=-\lambda_0\phi$. Then $\wp'(t,x,y):=\wp(t,x,y)\frac{\phi(y)}{\phi(x)}e^{-\lambda_0 t}$ satisfies $\int\wp'(t,x,y)dy=1$ thus defines a diffusion D with its Green function $G_D(x,y)=G_{\lambda_0}(x,y)\frac{\phi(y)}{\phi(x)}$. Suppose that there is a compact set K with non-empty interior such that $\int_K G_{\lambda_0}(x,y)\,d\mathrm{Vol}(y)=\infty$. It implies that $\int_K G_D(x,y)\,d\mathrm{Vol}(y)=\infty$. By the proof of Theorem 4.2.1.(ii) of [**Pi**], $G_{\lambda_0}(x,y)\frac{\phi(y)}{\phi(x)}e^{\lambda_0 t}=\infty$, which implies

$G_{\lambda_0}(x,y)=\infty$. By a theorem of Sullivan (Theorem 2.7 of [**S**]), there is a unique $(-\lambda_0)$-eigenfunction, thus $\phi(y)/\phi(x)$ is Γ-invariant. It follows that G_D is Γ-invariant thus by discretization (see the proof the main theorem of [**BL**]) there is a recurrent random walk μ_D on Γ with Green function G_D, which implies that Γ is virtually $\mathbb{Z},\mathbb{Z}^2$ or trivial [**V**]. ☐

Proposition 2.2. *We have, for $\lambda\in[0,\lambda_0)$, for any two points $x\neq y\in\widetilde{M}$:*

$$\frac{\partial^k}{\partial\lambda^k}G_\lambda(x,y) \;=\; k!\int_{\widetilde{M}^k}G_\lambda(x,x_1)G_\lambda(x_1,x_2)\cdots G_\lambda(x_k,y)\,d\,Vol^k(x_1,\cdots,x_k). \tag{2.2}$$

Proof. It follows directly from computation (See [**La**] for discrete case), e.g. for $k=1$,

$$\begin{aligned}\int_{\widetilde{M}}G_\lambda(x,z)G_\lambda(z,y)dz &= \int_0^\infty\int_0^\infty\int_{\widetilde{M}}e^{\lambda(t+u)}\wp(t,x,z)\wp(u,z,y)dzdtdu\\ &= \int_0^\infty\int_0^\infty e^{\lambda(t+u)}\wp(t+u,x,y)dtdu\\ &= \int_0^\infty\int_0^s e^{\lambda s}\wp(s,x,y)dtds=\int_0^\infty se^{\lambda s}\wp(s,x,y)ds=\frac{\partial}{\partial\lambda}G_\lambda(x,y).\end{aligned}$$

☐

Since the Green function is positive, by (2.2) for $k=1$ and 2, the map $\lambda\mapsto G_\lambda(x,y)$ is a convex increasing function. Since $G_\lambda(x,y)$ is analytic outside the spectrum as a resolvent, its derivative is finite as well, i.e.

$$\text{for all } \lambda<\lambda_0,\;\; \text{all } x\neq y\in\widetilde{M},\; \int_{\widetilde{M}}G_\lambda(x,z)G_\lambda(z,y)d\mathrm{Vol}(z)<+\infty. \tag{2.3}$$

For each $x\in\widetilde{M}$ and ξ in the geometric boundary $\partial\widetilde{M}$, there is a unique unit speed geodesic $\gamma_{x,\xi}$ starting from x in the equivalent class $[\xi]$ of ξ. The mapping $\xi\mapsto\dot{\gamma}_{x,\xi}(0)$ is a homeomorphism, which we denote by π_x^{-1}, from $\partial\widetilde{M}$ to the unit sphere $S_x\widetilde{M}$ in the tangent space at x to $\widetilde{M}$. Thus we will identify the unit tangent bundle $S\widetilde{M}$ with $\widetilde{M}\times\partial\widetilde{M}$.

For each $x\in\widetilde{M}$, $\partial\widetilde{M}$ is endowed with the Gromov metric

$$d_x(\xi,\eta)\;=\;e^{-a(\xi|\eta)_x},$$

where $0<a\leq 1$, is such that the sectional curvature κ satisfies $\kappa\leq -a^2$ on $\widetilde{M}$ and $(\xi|\eta)_x$ is the *Gromov product*

$$(\xi|\eta)_x=\lim_{y\to\xi,z\to\eta}\frac{1}{2}\left(d(x,y)+d(x,z)-d(y,z)\right). \tag{2.4}$$

The following properties follow from pinched negative curvature:

Proposition 2.3 ([**An1**]). *For all $\lambda\in[0,\lambda_0)$, every $\xi\in\partial\widetilde{M}$ there exist a positive $(-\lambda)$-eigenfunction $k_\lambda(x,y,\xi)$ of the Laplacian such that for each $x,y\in\widetilde{M}$,*

$$\lim_{z\to\xi}\frac{G_\lambda(y,z)}{G_\lambda(x,z)}\;=\;k_\lambda(x,y,\xi). \tag{2.5}$$

For any positive $(-\lambda)$*-eigenfunction* F*, there is a measure* ν_F *on* $\partial\widetilde{M}$ *such that*

$$F(y) \;=\; \int_{\partial\widetilde{M}} k_\lambda(x,y,\xi)\,d\nu_F(\xi).$$

Proposition 2.4 ([**H1**]). *Moreover, for all* $\lambda \in [0,\lambda_0)$*, there are constants* $\alpha(\lambda) > 0, C(\lambda) > 0$ *such that*

$$\frac{\|\nabla_y \ln k_\lambda(x,y,\xi) - \nabla_y \ln k_\lambda(x,y,\eta)\|}{(d_x(\xi,\eta))^\alpha} \;\le\; C(\lambda).$$

Proposition 2.5 ([**K1**]). *For three distinct points* $x, y, z \in \widetilde{M} \cup \partial\widetilde{M}$*, consider the function*

$$\theta^\lambda_x(y,z) \;:=\; \frac{G_\lambda(y,z)}{G_\lambda(y,x)G_\lambda(x,z)}. \tag{2.6}$$

There is a $\tau > 0$*, and a* τ*-Hölder continuous positive function* $\theta^\lambda_x(\xi,\eta)$ *on* $\partial\widetilde{M} \times \partial\widetilde{M} \backslash \Delta$ *such that*

$$\theta^\lambda_x(\xi,\eta) \;=\; \lim_{y\to\xi, z\to\eta} \theta^\lambda_x(y,z).$$

The function $\theta^\lambda_x(\xi,\eta)$, when it is finite as it is here, is called the *Naïm kernel* in potential theory [**N**]. Compare with the definition of the Gromov product (2.4).

Fix $x \in \widetilde{M}$. For $\xi \in \partial\widetilde{M}, y \in \widetilde{M}$, the Busemann function $\beta(x,y,\xi)$ is defined by

$$\beta(x,y,\xi) \;=\; \lim_{y_n\to\xi} d(y,y_n) - d(x,y_n). \tag{2.7}$$

As $\widetilde{M}$ is the universal cover of a closed manifold of negative curvature, we also use the thermodynamic formalism of the geodesic flow as in [**K1**], [**H1**], [**L2**].

The geodesic flow $g = \{g_t\}_{t\in\mathbb{R}}$ is defined on the unit tangent bundles SM and $S\widetilde{M}$. On SM, the geodesic flow is an Anosov flow. For a g-invariant probability measure m on SM, denote by $h_m(g)$ the measure-theoretic entropy of time-1 map g_1 with respect to m (see e.g. [**W**]) . For any continuous function φ, define the *topological pressure* $P(\varphi)$ *of* φ by

$$P(\varphi) \;:=\; \sup_m \left(h_m(g) + \int_{SM} \varphi\, dm \right), \tag{2.8}$$

where the supremum is taken over all g-invariant probability measures on SM.

Consider $v \in SM$. For a lift $\widetilde{v}$ in $S\widetilde{M}$, consider the geodesic $\gamma_{\widetilde{v}}(t)$ with initial tangent vector $\dot{\gamma_{\widetilde{v}}}(0) = \tilde{v}$. We will denote $\widetilde{v}^- = \gamma_{\widetilde{v}}(-\infty)$ and $\widetilde{v}^+ = \gamma_{\widetilde{v}}(+\infty)$. For all $\lambda \in [0,\lambda_0)$, the *potential function associated to* λ is the function on SM defined as

$$\varphi_\lambda(v) := -2\frac{d}{dt} \ln k_\lambda(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t), \widetilde{v}^+)\Big|_{t=0}.$$

We set $P(\lambda) := P(\varphi_\lambda)$ for $0 \le \lambda < \lambda_0$. Let m_λ to be the unique equilibrium probability measure of φ_λ, which attains the supremum in (2.8). It is mixing for the geodesic flow

g of M. The measure m_λ can be constructed from the *generalized family of Patterson-Sullivan measures associated to the potential function* φ_λ, characterized by the following proposition.

Proposition 2.6 ([**L2**]). *Fix* $x \in \widetilde{M}$. *For all* $\lambda \in [0, \lambda_0)$, *all* $y \in \widetilde{M}$, *there is a finite measure* μ_y^λ *on* $\partial\widetilde{M}$ *such that*

1) *the family* $y \mapsto \mu_y^\lambda$ *is* Γ*-equivariant:* $\mu_{\gamma y}^\lambda = \gamma_*(\mu_y^\lambda)$ *for* $\gamma \in \Gamma$ *and*
2) *for all* $\lambda \in [0, \lambda_0)$, μ_x^λ*-a.e.* $\xi \in \partial\widetilde{M}$,

$$\frac{d\mu_y^\lambda}{d\mu_x^\lambda}(\xi) \;=\; k_\lambda^2(x, y, \xi)e^{-P(\lambda)\beta(x,y,\xi)}.$$

The family is unique if we normalize by $\int_{M_0} \mu_y^\lambda(\partial\widetilde{M}) d\,Vol(y) = 1$.

Using Harnack inequality, the measures μ_x^λ defined above have finite total mass:

Proposition 2.7 (Harnack inequality [**CY**]). *There is a* $C_0 > 0$ *such that for all* $\lambda \in [0, \lambda_0]$, *for any positive function* f *which is* $(\Delta + \lambda)$*-harmonic on a domain* $\mathcal{D}$, *we have* $\|\nabla \ln f\|(x) \le \ln C_0$ *if* $d(x, \partial\mathcal{D}) > 1$.

Corollary 2.8. *There exists a constant* $C > 0$, *such that for all* $\lambda \in [0, \lambda_0)$, *all* $x \in \widetilde{M}$,

$$C^{-1} \le \mu_x^\lambda(\partial\widetilde{M}) \le C.$$

Proof. By Proposition 2.7 applied to $k_\lambda(x, y, \xi)$, for $x, y \in M_0$, $|\ln k_\lambda^2(x, y, \xi)|$ is bounded. By Proposition 2.7 again, the function φ_λ is bounded by $2\ln C_0$. It follows that the pressure $P(\lambda)$ is bounded. Thus, the Radon-Nikodym derivatives $\frac{d\mu_x^\lambda}{d\mu_y^\lambda}$ is bounded for $x, y \in M_0$ uniformly in λ. Since the total measure is 1, the corollary follows. □

Fix $x_0 \in \widetilde{M}$. By *Hopf parametrization*, i.e. by associating $(v^-, v^+, \beta(x_0, \gamma_v(0), v^+))$ to v, we identify $S\widetilde{M}$ with $(\partial\widetilde{M} \times \partial\widetilde{M} \backslash \Delta(\partial\widetilde{M})) \times \mathbb{R}$, where $\Delta(\partial\widetilde{M})$ is the diagonal embedding. Since $\theta_x^2(\xi, \eta)e^{2P(\lambda)(\xi|\eta)_x} d\mu_x(\xi)d\mu_x(\eta)$ is independent of x, we define a Γ-invariant, g_t-invariant measure $\widetilde{m}_\lambda$ by

$$d\widetilde{m}_\lambda(\xi, \eta, t) = \Omega_\lambda \theta_x^2(\xi, \eta)e^{2P(\lambda)(\xi|\eta)_x} d\mu_x^\lambda(\xi) \times d\mu_x^\lambda(\eta) \times dt \tag{2.9}$$

on $S\widetilde{M}$, which does not depend on x (it does not depend on x_0 either). Here, Ω_λ is the normalizing constant chosen so that the measure $\widetilde{m}_\lambda$ is equal to the Γ-invariant lift of the *probability* measure m_λ to $S\widetilde{M}$.

Remark 2.9. Note that we have a symmetric measure thanks to the fact that our potential function φ_λ is cohomologous to $\varphi_\lambda \circ \iota$ where ι is the flip map $v \mapsto -v$ (compare with asymmetric measure in [**PPS**] Section 3.7).

We can also identify the orthogonal two frame bundle $S^2\widetilde{M}$ with the triples of pairwise distinct points in $\partial\widetilde{M} \times \partial\widetilde{M} \times \partial\widetilde{M}$ by associating $(v, w \in v^\perp)$ to (v^+, v^-, w^+). The measure

$$d\tau_x^\lambda(a, b, c) := \Upsilon_\lambda \theta_x^\lambda(a, b)\theta_x^\lambda(b, c)\theta_x^\lambda(c, a)d\mu_x^\lambda(a)d\mu_x^\lambda(b)d\mu_x^\lambda(c) \tag{2.10}$$

does not depend on x and is Γ-invariant. Here Υ_λ is a normalizing constant so that the total measure satisfies

$$\tau_x^\lambda(S^2M) = 1. \tag{2.11}$$

Let us recall dynamical foliations of $T^1\widetilde{M}$ in order to define measures associated to μ_x^λ. For every $v \in T^1\widetilde{M}$, define the *strong stable manifold, strong unstable manifold, weak (or central) stable manifold and weak (or central) unstable manifold* of v as follows:

$$\begin{aligned} W^{ss}(v) &= \{w \in T^1\widetilde{M} : \lim_{t\to+\infty} d(g_t v, g_t w) = 0\}, \\ W^{uu}(v) &= \{w \in T^1\widetilde{M} : \lim_{t\to-\infty} d(g_t v, g_t w) = 0\}, \\ W^{cs}(v) &= \{w \in T^1\widetilde{M} : \exists s, \lim_{t\to+\infty} d(g_{t+s} v, g_t w) = 0\}, \\ W^{cu}(v) &= \{w \in T^1\widetilde{M} : \exists s, \lim_{t\to-\infty} d(g_{t+s} v, g_t w) = 0\}. \end{aligned}$$

The generalized Patterson-Sullivan measures μ_x^λ obtained in Proposition 2.6 induce measures on strong unstable manifolds and more generally any transversal to W^{cs} via the homeomorphism $(\pi_x^{-1})_*\mu_x^\lambda$: first, on $W^{uu}(v)$, we define

$$d\mu_\lambda^{uu}(w) = k^2(x_0, \pi w, w^+)e^{P(\lambda)\beta(\pi(w),x_0,w^+)}d\mu_{x_0}^\lambda(w),$$

which is independent of x_0 and is Γ-equivariant (see [**PPS**] Section 3.9 for example).

By extending the family μ_λ^{uu} on $W^{uu}(v)$, we define a family of measures μ_λ^{uu} on all transversals to the central stable manifolds on $S\widetilde{M} \simeq \widetilde{M} \times \partial\widetilde{M}$ in such a way that for two transversals through $\pi_x^{-1}(\xi)$ and $\pi_y^{-1}(\xi)$, respectively, the Radon-Nikodym derivative $\rho_\lambda(\pi_x^{-1}(\xi), \pi_y^{-1}(\xi))$ of the holonomy from $\pi_x^{-1}(\xi)$ to $\pi_y^{-1}(\xi)$ along the leaf $\widetilde{M} \times \{\xi\}$ is given by

$$\rho_\lambda(\pi_x^{-1}(\xi), \pi_y^{-1}(\xi)) = k_\lambda^2(x, y, \xi)e^{-P(\lambda)\beta(x,y,\xi)}.$$

Observe that μ_λ^{uu} on $S_x\widetilde{M}$ is $(\pi_x^{-1})_*\mu_x^\lambda$; note that

$$\frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) = e^{-tP(\lambda)}k_\lambda^2(\gamma_{\widetilde{v}}(t), \gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(\infty)),$$

and

$$\int_{S_{px}M} f(v)\, d\mu_\lambda^{uu}(v) = \int_{\partial\widetilde{M}} f(p \circ \pi_x^{-1}\xi) d\mu_x^{\lambda_0}(\xi), \tag{2.12}$$

$$\int_{S_{py}M} h(-u)\, d\mu_\lambda^{ss}(u) = \int_{\partial\widetilde{M}} h(p \circ \pi_y^{-1}\xi) d\mu_y^{\lambda_0}(\xi). \tag{2.13}$$

By a direct generalization of Margulis argument [**M1**] to Gibbs measures, one obtains the following proposition (see Section 4 for details and [**L1**] for the case $\lambda = 0$).

Proposition 2.10. *There exists a positive continuous function $D : (\widetilde{M} \times [0, \lambda_0)) \to \mathbb{R}_+$ such that*

$$\lim_{R\to\infty} e^{-RP(\lambda)} \int_{S(x,R)} G_\lambda^2(x, z)dz = D(x, \lambda).$$

Clearly, $x \mapsto D(x, \lambda)$ is Γ-invariant and depends only on $p(x) \in M$. The function $D(x, \lambda)$ will be described in Corollary 4.9.

Corollary 2.11. *For all $\lambda \in [0, \lambda_0)$, we have $P(\lambda) < 0$.*

Proof. Indeed, otherwise, we have by Proposition 2.7 and Proposition 2.10,

$$\int_{\widetilde{M}} G_\lambda(x,z)G_\lambda(z,y)d\mathrm{Vol}(z) \geq \frac{1}{C}\int_{1+d(x,y)}^{+\infty}\left(\int_{S(x,R)} G_\lambda^2(x,z)dz\right)dR \geq \frac{D(x,\lambda)}{C}\int_{1+d(x,y)}^{+\infty} dR.$$

The integral diverges, which is a contradiction with (2.3). □

The rest of the section is devoted to the proof of the following proposition due to Hamenstädt.

Proposition 2.12 ([**H3**], Corollary 5.5.1)**.** *There is a constant $C > 0$ such that for $R \geq 1$,*

$$\int_{S(x,R)} G_{\lambda_0}^2(x,z)dz \leq C.$$

Proof. (cf. [**G1**], Lemma 2.5. We repeat, with our notations, the proof in [**H3**]. One reason is the comfort of the reader.) It follows from Harnack inequality (Proposition 2.7)(see [**An2**] Remark in the page 94) that there exists a constant C_0' such that for $0 \leq \lambda \leq \lambda_0$ and x, y, z such that $d(x,z) \geq 1, d(x,y) \geq 1$, we have

$$G_\lambda(x,z)G_\lambda(x,y) \leq C_0' G_\lambda(z,y).$$

Thus for any $\lambda \in [0, \lambda_0]$, any $x, z \in \widetilde{M}$ and any $y \in \widetilde{M}$ with $d(x,z) \geq 1, d(x,y) \geq 1$,

$$G_\lambda(x,z) = \frac{G_\lambda(x,z)G_\lambda(x,y)}{G_\lambda(x,y)} \leq C_0' \frac{G_\lambda(z,y)}{G_\lambda(x,y)},$$

Letting y go to $\xi \in \partial\widetilde{M}$, we obtain

$$G_\lambda^2(x,z)e^{-P(\lambda)d(x,z)} \leq {C_0'}^2 k_\lambda^2(x,z,\xi)e^{-P(\lambda)d(x,z)}.$$

A transversal T to the central stable manifold W^{cs} is said to be *ε-transversal* if the maximum angle between T and the strong unstable manifold W^{uu} is smaller than ε. Observe that $g_R(S_xM)$ is ε-transversal to W^{cs} for $R \geq R_0$ for some $R_0 > 0$.

Lemma 2.13. *Let T be ε-transversal. There is C_ε such that for all $w \in T$, $\lambda \in [0, \lambda_0]$,*

$$\mu_\lambda^{uu}(B(w,1) \cap T) \geq C_\varepsilon^{-1}.$$

Proof. Let $B = B(w,1) \cap T$. By the minimality of W^{cs} on SM, there is r large enough that $\cup_{b \in B} B^{cs}(b,r)$ covers at least one fundamental domain of SM, where $B^{cs}(b,r)$ is the ball of center b and radius r in W^{cs}. By Harnack inequality, $\rho_\lambda(b,z)$ is bounded from above on $B^{cs}(b,r)$ uniformly in λ and b. Therefore there is C_ε such that for all λ

$$\int_B \int_{B^{cs}(b,r)} \rho_\lambda(b,z)d\mathrm{Vol}(z)d\mu_\lambda^{uu}(b) \leq C_\varepsilon \mu_\lambda^{uu}(B).$$

On the other hand, by our choice of normalization in Proposition 2.6,

$$\int_B \int_{B^{cs}(b,r)} \rho_\lambda(b,z)d\mathrm{Vol}(z)d\mu_\lambda^{uu}(b) \geq \int_{M_0}\int_{\partial\widetilde{M}} \frac{d\mu_x}{d\mu_{x_0}}(\xi)d\mu_{x_0}(\xi)dm(x) = 1.$$

□

Using all the above estimates, for any $R \geq R_0$, $\lambda < \lambda_0$ and a ball $B(w,1)$ of radius 1 in $S(x,R)$,

$$\begin{aligned}
\int_{B(w,1)} G_\lambda^2(x,z)e^{-P(\lambda)R}dz &\leq C_0^2G_\lambda^2(x,w)e^{-P(\lambda)R} \leq C_0^2{C_0'}^2\frac{dg_R\mu_\lambda^{uu}}{d\mu_\lambda^{uu}}(w) \\
&\leq \frac{C_0^4{C_0'}^2}{\mu_\lambda^{uu}(B(w,1))}\int_{B(w,1)}\frac{dg_R\mu_\lambda^{uu}}{d\mu_\lambda^{uu}}(z)d\mu_\lambda^{uu}(z) \\
&\leq C_0^4{C_0'}^2\frac{g_R\mu_\lambda^{uu}(B(w,1))}{\mu_\lambda^{uu}B(w,1)} \\
&\leq C_\varepsilon C_0^6{C_0'}^2\mu_\lambda^{uu}g_{-R}(B(w,1)).
\end{aligned}$$

By covering $S(x,R)$ by balls of radius 1, the intersection of which is of bounded multiplicity, we obtain a constant C such that for all $\lambda < \lambda_0$ and for $R > R_0$,

$$\int_{S(x,R)} G_\lambda^2(x,z)e^{-P(\lambda)R}dz \leq C. \tag{2.14}$$

Since $P(\lambda) < 0$ for all $\lambda < \lambda_0$, there is a constant $C > 0$ such that for all $\lambda \in [0,\lambda_0)$, all $x \in \widetilde{M}$, all $R \geq 1$,

$$\int_{S(x,R)} G_\lambda^2(x,z)dz \leq C.$$

Proposition 2.12 follows by letting λ go to λ_0. □

Corollary 2.14. *For $T > 0$, $P_T(\lambda)$ be the Pressure of the function $\frac{T}{2}\varphi_\lambda$. Then there exists a constant $C(T)$ and R_0 such that for all $\lambda \in [0,\lambda_0), R \geq R_0, x \in \widetilde{M}$,*

$$e^{-RP_T(\lambda)}\int_{S(x,R)} G_\lambda^T(x,z)dz \ \leq C(T).$$

Proof. We have as above

$$G_\lambda^T(x,z)e^{-P_T(\lambda)d(x,z)} \leq {C_0'}^T k_\lambda^T(x,z,\xi)e^{-P_T(\lambda)d(x,z)}.$$

We can also apply Proposition 2.6 to the Hölder continuous function $\frac{T}{2}\varphi_\lambda$ instead of φ_λ. We obtain a family of measures $\mu_x^{\lambda,T}$ on $\partial\widetilde{M}$ such that for all $\lambda \in [0,\lambda_0)$, μ_x^λ-a.e. $\xi \in \partial\widetilde{M}$,

$$\frac{d\mu_y^{\lambda,T}}{d\mu_x^{\lambda,T}}(\xi) \ = \ k_\lambda^T(x,y,\xi)e^{-P_T(\lambda)\beta(x,y,\xi)}$$

and $\int_{M_0} \mu_y^{\lambda,T}(\partial\widetilde{M})d\mathrm{Vol}(y) = 1$. We can therefore associate measures $\mu^{uu}_{\lambda,T}$ on transversals to the central stable manifolds such that the holonomy from $\pi_x^{-1}(\xi)$ to $\pi_y^{-1}(\xi)$ along the leaf $\widetilde{M} \times \{\xi\}$ is given by

$$\rho_\lambda(\pi_x^{-1}(\xi), \pi_y^{-1}(\xi)) = k_\lambda^T(x,y,\xi)e^{-P_T(\lambda)\beta(x,y,\xi)}.$$

The same computation yields the analog of (2.14). □

Corollary 2.15. *Given ε, there exists a constant C_2 with the following property: let A be a $(m-1)$-dimensional submanifold of $\widetilde{M}$ with bounded geometry, $x \in \widetilde{M}, d(x,A) \geq 1$, and assume that the lift $\widetilde{A}^x := \{(y, -v_y^x); y \in A\}$ to $S\widetilde{M}$ is ε transversal to W^{cs}, then*

$$\int_A G^2_{\lambda_0}(x,y)\, dy \ \leq \ C_2.$$

Here, v_y^x is the unit vector based on y pointing towards x.

Proof. Let $y \in A$ and let $B(y,1)$ be the ball of radius 1 about y in A. We obtain as above, uniformly in $\lambda \in [0, \lambda_0)$,

$$\int_{B(y,1)} G^2_\lambda(x,w)e^{-P(\lambda)d(x,w)}\, dw \ \leq \ CC_\varepsilon \mu^{uu}(B_y),$$

where B_y is the subset of $S_x\widetilde{M}$ defined by $B_y := \{\dot\gamma_{v_w^x}(d(w,x)) : w \in B(y,1)\}$. By bounded geometry, we can cover A by balls $B(y_i, 1)$ with bounded multiplicity. The subsets B_{y_i} cover a part of $S_x\widetilde{M}$ with the same bounded multiplicity; therefore we find a constant C_2 such that, for all $\lambda \in [0, \lambda_0)$, all x, A such that $\widetilde{A}$ is ε transversal to W^{cs},

$$\int_A G^2_\lambda(x,y)e^{-P(\lambda)d(x,y)}\, dy \ \leq \ C_2.$$

Since $P(\lambda) < 0$ for all $\lambda < \lambda_0$, we have $\int_A G^2_\lambda(x,y)\, dy \ \leq \ C_2$. The corollary follows by letting $\lambda \to \lambda_0$ and the Monotone Convergence Theorem. □

3. Ancona-Gouëzel inequality

Definition 3.1. *Let $v \in S\widetilde{M}$. The cone $\mathcal{C}(v)$ based on v is defined by:*

$$\mathcal{C}(v) \ := \ \{y; y \in \widetilde{M}, \angle_x(v,y) \leq \pi/2\},$$

where $\angle_x(v,y)$ denotes the angle between $v \in T_x\widetilde{M}$ and the geodesic going from x to y.

We denote $\partial\mathcal{C}(v) := \{y; y \in \widetilde{M}, \angle_x(v,y) = \pi/2\}$, Observe that $\widetilde{M} = \mathcal{C}(v) \cup \mathcal{C}(-v)$ and $\partial\mathcal{C}(v) = \mathcal{C}(v) \cap \mathcal{C}(-v)$.

Let $\mathcal{D}$ be a open subset of $\widetilde{M}$. For $x, y \in \mathcal{D}$, the relative Green function $G_\lambda(x, y : \mathcal{D})$ is the Green function of the process killed when entering $\widetilde{M} \setminus \mathcal{D}$. It can be defined as the Green function of the operator $\Delta + \lambda$ on $\mathcal{D}$ with Dirichlet conditions on the boundary. Clearly, for all set $\mathcal{D}, \mathcal{C}, \mathcal{C} \subset \mathcal{D}$ all $\lambda \leq \lambda_0$, and all $x, y \in \mathcal{C}$, $G_\lambda(x, y : \mathcal{C}) \leq G_\lambda(x, y : \mathcal{D})$.

In general, if A is a closed subset of the open $\mathcal{D}$, and $x \in \mathcal{D} \setminus A$, there is a measure ϖ_x^λ on A such that:

$$G_\lambda(x, z : \mathcal{D}) \;=\; \int_A G_\lambda(y, z : \mathcal{D})\, d\varpi_x^\lambda(y) + G_\lambda(x, z : \mathcal{D} \setminus A). \tag{3.1}$$

We recall the proof in Section 8. Assume that A is a $(n-1)$-dimensional submanifold of $\widetilde{M}$, with bounded geometry. Then, the measure ϖ_x^λ has a density with respect to the induced Lebesgue measure dy on A. We can estimate some regularity of this density with some geometric hypotheses:

Proposition 3.2. *Let A be a $(n-1)$ dimensional submanifold of $\mathcal{D}$ with bounded geometry. Set A_1 for the set of points of A at distance at least 1 from $\mathcal{D}^c$. There exists a constant C_3 such that for $\lambda \in [0, \lambda_0]$, for any positive function F on A_1, any $x \in \mathcal{D}$ with $d(x, \mathcal{D}) > 1$,*

$$\int_{A_1} F(y) d\varpi_x^\lambda(y) \le\; C_3 L(F)^2 \int_A G_\lambda(x, y) F(y)\, dy,$$

where $L(F) := e^{\sup_A \|\nabla \ln F\|}$ is the (multiplicative) Lipschitz constant of F.

Proposition 3.3. *Let $\mathcal{C}$ be an open domain, $\mathcal{C} \subset \mathcal{D}, d(\mathcal{C}, \partial\mathcal{D}) > 1$. Let $x \in \mathcal{C}, A := \partial\mathcal{C}$ and let ϖ_x^λ be the distribution in (3.1) on A. There exists a constant C_3 such that if $d(x, A) > 1$, then for $\lambda \in [0, \lambda_0]$, for any positive function F on A_1,*

$$C_3^{-1}(L(F))^{-2} \int_A G_\lambda(x, y : \mathcal{D}) F(y)\, dy \;\le\; \int_A F(y) d\varpi_x^\lambda(y).$$

We chose the same constant C_3 in both Propositions for convenience. A priori, the constant C_3 depends on the geometry of $\widetilde{M}$ and A. Since we use Harnack inequality on $\widetilde{M}$ in the proof, it might also deteriorate when $\lambda \to -\infty$. For the comfort of the reader, probabilistic proofs are recalled in Section 8.

3.1. **Ancona-Gouëzel inequality.**

The key property of the λ-Green functions for $0 \le \lambda \le \lambda_0$ is the following *uniform* Ancona inequality, which we call *Ancona-Gouëzel inequality.*

Theorem 3.4. *With the above notations, there are constants C_4, R_0 such that for all $\lambda \in [0, \lambda_0]$, all points (x, y, z) such that y is on the geodesic segment $[xz]$ from x to z and $d(x, y) \ge R_0, d(y, z) \ge R_0$,*

$$C_4^{-1} G_\lambda(x, y : \mathcal{D}) G_\lambda(y, z : \mathcal{D}) \;\le\; G_\lambda(x, z : \mathcal{D}) \;\le\; C_4 G_\lambda(x, y : \mathcal{D}) G_\lambda(y, z : \mathcal{D}) \tag{3.2}$$

for all open set $\mathcal{D}$ containing $\mathcal{C}(g_{-1}v) \cup \mathcal{C}(-g_{d(x,z)+1}v)$, where $v \in S_x\widetilde{M}$ is the initial vector of the geodesic $[xz]$.

Theorem 3.4 was proven by A. Ancona for $\lambda < \lambda_0$ ([**An1**]). The first inequality in (3.2) follows from Harnack inequality uniformly in $0 \le \lambda \le \lambda_0$ (again, see [**An2**], page 94). The new fact here is that the second inequality (3.2) holds when $\lambda = \lambda_0$ as well, with the same constant C_4, so that the consequences of Theorem 3.4 are now uniform in $\lambda \in [0, \lambda_0]$ as in [**G1**]. The Ancona inequality follows from the pre-Ancona inequality in the following Proposition.

Proposition 3.5. *Let x, y, z be points on a geodesic γ in this order, v the tangent vector to γ at x. Then, there exists $\varepsilon > 0, R_2$ such that if $r \geq R_2$ and $d(x,y) > r+1, d(y,z) > r+1$,*

$$G_{\lambda_0}(x, z : B(y,r)^c \cap \mathcal{C}(g_{-1}v) \cap \mathcal{C}(-g_{d(x,z)+1}v)) \leq 2^{-e^{\varepsilon r}}.$$

Proof. As in [**G1**], we will construct $N = e^{\varepsilon r}$ barriers, for a positive constant ε which we will specify as follows.



Figure 1. Ancona-Gouëzel inequality

For $i = 1, \cdots, N$, let $X_i = ((N+2i-1)\pi/4N, (N+2i)\pi/4N) \subset [\pi/4, 3\pi/4]$. Choose θ_i from X_i, for $i = 1, \cdots, N$. The circumference of a great circle of a ball $B(y, r-1)$ is at least $e^{a_0(r-1)}$, where a_0 is a geometric constant coming from the fact that the curvature is negative and bounded away from 0. Thus the intersections $\{A_i\}$'s of $B(y, r-1)^c$ and the cones $\{w : \angle_y(x,w) = \theta_i\}$ of angle θ_i at y, are of distance between them bounded below by $e^{(a_0-\varepsilon)r}$ for all r large enough. Set $\mathcal{D} := B(y,r)^c \cap \mathcal{C}(g_{-1}v) \cap \mathcal{C}(-g_{d(x,z)+1}v)$. Each set $A_i \cap \mathcal{D}$ separate $\mathcal{D}$ into two disjoint open sets. Let $\mathcal{C}_i$ be the one containing x. Then $z \notin \mathcal{C}_i$. Moreover, the sets $A_i \cap \mathcal{D} = \partial\mathcal{C}_i$ have uniform bounded geometry and do not intersect $\partial\mathcal{C}(g_{-1}v) \cup \mathcal{C}(-g_{d(x,z)+1}v)$ (see Figure 1).

By (3.1), we may write:

$$\begin{aligned}
G_{\lambda_0}(x,z:\mathcal{D}) &= \int_{A_1\cap\mathcal{D}} G_{\lambda_0}(u_1,z:\mathcal{D})\,d\varpi_x^\lambda(u_1)\\
&= \int_{A_1\cap\mathcal{D}}\int_{A_2\cap\mathcal{D}} G_{\lambda_0}(u_2,z:\mathcal{D})\,d\varpi_x^\lambda(u_1)d\varpi_{u_1}^\lambda(u_2)\\
&= \int_{A_1\cap\mathcal{D}}\cdots\int_{A_N\cap\mathcal{D}} G_{\lambda_0}(u_N,z:\mathcal{D})\,d\varpi_x^\lambda(u_1)\cdots d\varpi_{u_{N-1}}^\lambda(u_N)\\
&\le \int_{A_1\cap\mathcal{D}}\cdots\int_{A_N\cap\mathcal{D}} G_{\lambda_0}(u_N,z)\,d\varpi_x^\lambda(u_1)\cdots d\varpi_{u_{N-1}}^\lambda(u_N)
\end{aligned}$$

where ϖ_u^j is the distribution on $A_j\cap\mathcal{D}$ given by (3.1). (Observe that $G_\lambda(u_j,z):\mathcal{D}-A_j)=0$ since u_i,z are separated by A_j. Observe that, by Proposition 2.7, for all $u_N\in A_N, ||\nabla_{u_N}\ln G_{\lambda_0}(u_N,z)||\le \ln C_0$. By construction, $d(A_N\cap\mathcal{D},B(y,r-1))\ge 1$ and for all $u_{N-1}\in A_{N-1}, d(u_{N-1},B(y,r-1))\ge 1$. So, we may apply Proposition 3.2 and obtain a constant $C_5=C_3C_0^2$ such that

$$\int_{A_N\cap\mathcal{D}} G_{\lambda_0}(u_N,z)\,d\varpi'_{u_{N-1}}(u_N)\ \le\ C_5\int_{A_N} G_{\lambda_0}(u_{N-1},u_N)G_{\lambda_0}(u_N,z)\,du_N,$$

where ϖ'_z is the distribution on $\overline{A_N\cap\mathcal{D}}$ associated with (3.1) for the domain $B(y,r-1)^c\cap\mathcal{C}(g_{-1}v)\cap\mathcal{C}(-g_{d(x,z)+1}v)$. Since $\mathcal{D}\subset B(y,r-1)^c\cap\mathcal{C}(g_{-1}v)\cap\mathcal{C}(-g_{d(x,z)+1}v)$, we have $\varpi_z^N\le\varpi'_z$ on $A_N\cap\mathcal{D}$ and therefore

$$\int_{A_N\cap\mathcal{D}} G_{\lambda_0}(u_N,z)\,d\varpi^N_{u_{N-1}}(u_N)\ \le\ C_5\int_{A_N} G_{\lambda_0}(u_{N-1},u_N)G_{\lambda_0}(u_N,z)\,du_N.$$

The right hand side satisfies for all $u_{N-1}\in A_{N-1}$,

$$||\nabla_{u_{N-1}}\int_{A_N} G_{\lambda_0}(u_{N-1},u_N)G_{\lambda_0}(u_N,z)\,du_N||\le C_0\int_{A_N} G_{\lambda_0}(u_{N-1},u_N)G_{\lambda_0}(u_N,z)\,du_N$$

because it is an integral in the variable u_N of the functions $G_{\lambda_0}(u_{N-1},u_N)$ with that property. We can iterate the application of Proposition 3.2 and obtain

$$\begin{aligned}
G_{\lambda_0}(x,z:\mathcal{D}) &\le C_5^N\int_{A_1}\cdots\int_{A_N} G_{\lambda_0}(x,u_1)G_{\lambda_0}(u_1,u_2)\cdots G_{\lambda_0}(u_N,z)\,du_1\cdots du_N\\
&= C_5^N\,\langle\delta_x,L_1\cdots L_N G_{\lambda_0}(u_n,z)\rangle\le C_5^N||L_1\cdots L_N G_{\lambda_0}(u_n,z)||\\
&\le C_5^N\prod||L_i||\left(\int_{A_N} G_{\lambda_0}^2(u_N,z)du_N\right)^{1/2},
\end{aligned}$$

where $L_i:L^2(A_{i+1})\to L^2(A_i)$ is defined by $L_if(u_i)=\int G_{\lambda_0}(u_i,u_{i+1})f(u_{i+1})du_{i+1}$.

Observe that there is C such that, for all $z,d(z,A_N)\ge 1$, $\int_{A_N} G_{\lambda_0}^2(u_N,z)du_N\le C$ by Corollary 2.15 as we have the needed tranversality for $\widetilde{A}_N^z$. Thus it suffices to show that there exist $\theta_1,\cdots,\theta_N$ for which $f_i(\theta_1,\cdots,\theta_N)=||L_i||^2\le\int\int G_{\lambda_0}^2(u_i,u_{i+1})du_{i+1}du_i<\frac{1}{4C_5}$ for all i. Now choose θ_i uniformly from X_i. We claim that the expectation of f_i

with respect to normalized measures $\frac{4}{\pi}Nd\theta_i$ satisfies

$$\mathbb{E}(f_i) \leq \frac{e^{-\varepsilon r}}{5C_5},$$

if ε is small enough. It would imply that $\mathbb{E}(\sum f_i) \leq \frac{Ne^{-\varepsilon r}}{5C_5} < \frac{1}{4C_5}$, which would in turn imply that $\sum f_i(\theta_1, \cdots, \theta_N) < \frac{1}{4C_5}$ for some $\{\theta_1, \cdots, \theta_N\}$, thus $f_i(\theta_1, \cdots, \theta_N) < \frac{1}{4C_5}$ for all i.

Now it remains to prove the claim. For $a \in A_i, b \in A_{i+1}$, let $\Phi(a, b, \theta_i, \theta_{i+1}) = \gamma_0^{-1}\gamma_1 \in \Gamma$ where γ_0 and γ_1 are the first elements of Γ in the lexicographical order such that

$$d(\gamma_0 y, a) < \text{diam } M \text{ and } d(\gamma_1 y, b) < \text{diam } M.$$

Denoting by $d\mu(a, b, \theta_i, \theta_{i+1})$ the product of the Lebesgue measures on A_i, A_{i+1} and of $\frac{16}{\pi^2}N^2 d\theta_i d\theta_{i+1}$, we have

$$\begin{aligned}
\mathbb{E}(f_i) &= \int G^2_{\lambda_0}(a, b) d\mu(a, b, \theta_i, \theta_{i+1}) \\
&\leq C_0^{2\text{diam}M} \sum_{\gamma\in\Gamma} G^2_{\lambda_0}(y, \gamma y)\mu(\{(a, b, \theta_i, \theta_{i+1}) : \Phi(a, b, \theta_i, \theta_{i+1}) = \gamma\}) \\
&\leq C_0^{2\text{diam}M} \int_{\widetilde{M}} G^2_{\lambda_0}(y, w) A(w) d\text{Vol}(w),
\end{aligned}$$

where $A(z) = \mu(\{(a, b, \theta_i, \theta_{i+1}) : z \in \Phi(a, b, \theta_i, \theta_{i+1})M_0\})/\text{vol}(M)$ and $C_0^{2\text{diam}M}$ comes from Proposition 2.7. (Here M_0 is a fundamental domain containing y.) The first inequality holds since

$$G_{\lambda_0}(a, b) = G_{\lambda_0}(\gamma_0^{-1}a, \gamma_0^{-1}b) \leq C_0^{2\text{diam}M} G_{\lambda_0}(y, \gamma_0^{-1}\gamma_1 y).$$

Observe that the angles $\angle_y(\gamma_0 y, a), \angle_y(\gamma_1 y, b)$ are at most $e^{-a_0 r}$. Let us now estimate $A(w)$ for $w \in \widetilde{M}$. If ε is chosen small enough, this implies that $\angle_y(\gamma_0 y, \gamma_1 y) \geq 1/2e^{-\varepsilon r}$, so that γ_0 and γ_1 have at most $Ce^{a_1\varepsilon r}$ letters in common, for some constants a_1 and C depending on the bounds of the curvature and of the inner diameter of the fundamental domain M_0. Therefore, for a fixed $\gamma \in G$, there are at most $|\gamma|Ce^{a_1\varepsilon r}$ possible values of γ_0. For fixed γ, γ_0,

$$\begin{aligned}
\mu\{(a, b, \theta_i, \theta_{i+1}) &: \Phi(a, b, \theta_i, \theta_{i+1}) = \gamma, d(a, \gamma_0 y) \leq \text{diam } M, d(b, \gamma_0\gamma y) \leq \text{diam } M\} \\
&\leq CN^2 e^{-a_0(d(y,a)+d(y,b))} \leq CN^2 e^{-a_0|\gamma|}.
\end{aligned}$$

Since word metric on Γ and the hyperbolic metric are quasi-isometric, $|\gamma| \geq a_1 d(y, w) - 2$. Thus

$$A(w) \leq d(y, w)Ce^{a_1\varepsilon r}e^{2\varepsilon r}e^{-a_0 a_1 d(y,w)}.$$

It follows that there exists R_2 such that if ε is chosen small enough and $r \geq R_2$,

$$\begin{aligned}
\mathbb{E}(f_i) &\leq Ce^{(2+a_1)\varepsilon r} \int_r^\infty Re^{-a_0 a_1 R} \int_{S(y,R)} G^2_{\lambda_0}(y, z) dR \\
&\leq Ce^{(2+a_1)\varepsilon r} \int_r^\infty Re^{-a_0 a_1 R} dR \leq Ce^{((3+a_1)\varepsilon - a_0 a_1)r} < \frac{e^{-\varepsilon r}}{5C_5},
\end{aligned}$$

where we used Proposition 2.12 for the second inequality. □

Theorem 3.4 follows from Proposition 3.5 by an inductive argument (see also [**G1**], [**GL**]). Indeed, let $x, y, z, \mathcal{D}$ be as in Theorem 3.4, $\lambda \in [0, \lambda_0]$. We want to estimate from above

$$\frac{G_\lambda(x,z:\mathcal{D})}{G_\lambda(x,y:\mathcal{D})G_\lambda(y,z:\mathcal{D})}.$$

Set $\Psi(\alpha,\beta)$ the highest possible value of this ratio for $x, y, z, \mathcal{D}$ as in Theorem 3.4, $d(x,y) \leq \alpha, d(y,z) \leq \beta$. Assume without loss of generality that $\alpha \geq \beta$. We claim that there is $\theta < 1$ and R such that, if $\alpha, \beta \geq R$,

$$\Psi(\alpha,\beta) \;\leq\; (1+\theta^\alpha)\Psi(\alpha/2,\beta). \tag{3.3}$$

It follows that for all (α,β),

$$\Psi(\alpha,\beta) \;\leq \Pi_{k\in\mathbb{N}}(1+\theta^{2^k R})^2\Psi(2R,2R).$$

This shows Theorem 3.4 since the infinite product is converging and $\Psi(2R,2R)$ is finite. It remains to prove the claim. Consider $(x,y,z,\mathcal{D})$ as in Theorem 3.4, $d(x,y) \leq \alpha, d(y,z) \leq \beta$ achieving the maximum $\Psi(\alpha,\beta)$. Let x' be the point in the segment $[x,y]$ with $d(x',y) = 0.4\,\alpha$. Using (3.1) with the sphere $S(x', 0.1\alpha)$ of points at distance 0.1α from x', we see that, with $w = y$ or z, we can write, if $\alpha > 10R_2$,

$$G_\lambda(x,w:\mathcal{D}) \;=\; \int_{S(x',0.1\alpha)} G_\lambda(w',w:\mathcal{D})\,d\varpi_x^\lambda(w') + G_\lambda(x,w:\mathcal{D}\cap B(x',0.1\alpha)^c). \tag{3.4}$$

By Proposition 3.5, for all $\lambda, 0 \leq \lambda \leq \lambda_0$,

$$G_\lambda(x,w:\mathcal{D}\cap B(x',0.1\,\alpha)^c) \leq G_{\lambda_0}(x,w:\mathcal{D}\cap B(x',0.1\,\alpha)^c) \leq 2^{e^{-(0.1\,\alpha)}}$$

for $w = y$ or z whereas, by Proposition 3.3 and Harnack inequality (Proposition 2.7) there is a positive c such that, for all $\lambda, 0 \leq \lambda \leq \lambda_0$, the first term is at least c^α. If α is large enough, it will suffice to control the first term of (3.4).

Consider for any $w' \in S(x', 0.1\,\alpha)$, the point $y(w')$ on the geodesic $[w', z]$ at distance β from z and the point $x'(w')$ on the geodesic $[x,z]$ at distance $d(w',z)$ from z. Clearly,

$$G_\lambda(w',z:\mathcal{D}) \;\leq\; \Psi(0.5\,\alpha,\beta)G_\lambda(w',y(w'):\mathcal{D})G_\lambda(y(w'),z:\mathcal{D}).$$

Since $d(w',z) \geq \beta + 0.3\,\alpha$, there is a constant a_0 depending only on the curvature bounds such that

$$d(y,y(w')) \;\leq\; e^{-(0.3\,a_0\alpha)}d(x'(w'),w') \;\leq\; e^{-(0.3\,a_0\alpha)}0.1\,\alpha.$$

Choosing $\theta_0 < 1$ such that for α large enough, $C_0^{\left(0.6\,\alpha e^{-0.3\,a_0\alpha)}\right)} \leq 1 + \theta_0^\alpha$, we get by Proposition 2.7 that for all $w' \in S(x', 0.1\,\alpha)$,

$$G_\lambda(w',y(w'):\mathcal{D})G_\lambda(y(w'),z:\mathcal{D}) \leq (1+\theta_0^\alpha)G_\lambda(w',y:\mathcal{D})G_\lambda(y,z:\mathcal{D}).$$

Reporting in (3.4) with $w = z$, we obtain, if α is large enough:

$$\begin{aligned} G_\lambda(x,z:\mathcal{D}) &= \int_{S(x',0.1\,\alpha)} G_\lambda(w',z:\mathcal{D})\,d\varpi_x^\lambda(w')(1+o(\theta_0^\alpha)) \\ &\leq (1+\theta_0^\alpha)\Psi(0.5\,\alpha,\beta)G_\lambda(y,z:\mathcal{D})\int_{S(x',0.1\,\alpha)} G_\lambda(w',y:\mathcal{D})\,d\varpi_x^\lambda(w')(1+o(\theta_0^\alpha)) \\ &= (1+\theta_0^\alpha)(1+o(\theta_0^\alpha))\Psi(\alpha/2,\beta)G_\lambda(y,z:\mathcal{D})G_\lambda(x,y:\mathcal{D}), \end{aligned}$$

where we used (3.4) with $w = y$ in the last line. This achieves the proof of the claim and of Theorem 3.4.

We use the following notation throughout this article: $\sim^a$ means that the ratios between the two sides are bounded by a.

Corollary 3.6. *There are constants C_8, R_1 such that, for all $\lambda \in [0, \lambda_0]$, all $v \in \widetilde{M}$, all $y, y' \notin \mathcal{C}(g_{-R_1}v)$ and all $z \in \mathcal{C}(g_{R_1}v)$,*

$$G_\lambda(y,z) \ \sim^{C_8}\ G_\lambda(y, \gamma_v(0)) G_\lambda(\gamma_v(0), z), \qquad \frac{G_\lambda(y,z)}{G_\lambda(y',z)} \sim^{C_8^2} \frac{G_\lambda(y,\gamma_v(0))}{G_\lambda(y',\gamma_v(0))}. \tag{3.5}$$

Proof. Let $y \notin \mathcal{C}(g_{-R}v), z \in \mathcal{C}(g_R v)$. If R is large enough, on the geodesic $[yz]$, the closest point $w(y,z)$ to $\gamma_v(0)$ satisfies $d(w(y,z), \gamma_v(0)) \leq 1$. The first inequality in (3.5) follows directly from (3.2) and Proposition 2.7, the second from the first applied to $y, y' \notin \mathcal{C}(g_{-R_1}v)$. □

3.2. **Martin boundary.** We now follow Section 6 of [**AnS**] simultaneously for all $\lambda \in [0, \lambda_0]$ to obtain Propositions 2.3, 2.4, 2.5 uniformly in $\lambda \leq \lambda_0$. For $x, y, z \in \widetilde{M}, \lambda \in [0, \lambda_0]$, set

$$k_\lambda(x,y,z) \ := \ \frac{G_\lambda(y,z)}{G_\lambda(x,z)}.$$

The function $k_\lambda(x,y,z)$ is clearly $(\Delta + \lambda)$-harmonic in y on $\widetilde{M} \setminus \{z\}$.

Lemma 3.7. *There are constants $C > 1, K < 1$ such that for all geodesic γ and all $x, y \notin \mathcal{C}(\dot\gamma(-2R_1 - T)), z, w \in \mathcal{C}(\dot\gamma(2R_1)), \lambda \in [0, \lambda_0], T > 0$,*

$$\Big| \ln \frac{k_\lambda(x,y,z)}{k_\lambda(x,y,w)} \Big| \ \leq\ C K^T.$$

Proof. It suffices to prove the case $T = 2nR_1$ for $n \in \mathbb{N}$. For $v \in S\widetilde{M}$, denote $\mathcal{C}_{\pm 1}(v) := \mathcal{C}(g_{-1}v) \cap \mathcal{C}(-g_1(v))$. Fix a geodesic γ and points $z, w \in \mathcal{C}(\dot\gamma(2R_1))$. for $x, y \in \mathcal{C}_{\pm 1}(\dot\gamma(-2nR_1))$, denote

$$k_\lambda(x,y,z;n) = \frac{G_\lambda(y, z : \mathcal{C}(\dot\gamma(-2nR_1 - 2)))}{G_\lambda(x, z : \mathcal{C}(\dot\gamma(-2nR_1 - 2)))}.$$

The following numbers $\overline{\theta}(n), \underline{\theta}(n)$ are well defined for $n \in \mathbb{N}$ and a priori smaller than C_8^4 (and larger than $(C_8^4)^{-1}$) independently of $\lambda \in [0, \lambda_0]$, the geodesic γ and $z, w \in \mathcal{C}(\dot\gamma(2R_1))$ by (3.5) :

$$\overline{\theta}(n) \ := \sup_{x,y \in \mathcal{C}_{\pm 1}(\dot\gamma(-2nR_1))} \frac{k_\lambda(x,y,z;n)}{k_\lambda(x,y,w;n)} \qquad \underline{\theta}(n) \ := \inf_{x,y \in \mathcal{C}_{\pm 1}(\dot\gamma(-2nR_1))} \frac{k_\lambda(x,y,z;n)}{k_\lambda(x,y,w;n)}.$$

Let $x, y \in \mathcal{C}_{\pm 1}(-2(n+1)R_1))$. We apply Proposition 8.1 with $\mathcal{D} = \widetilde{M}$ and the separating $A = \partial\mathcal{C}(\dot\gamma(-2nR_1))$. Denote $\varpi_x^\lambda, \varpi_y^\lambda$ the hitting distributions on $\partial\mathcal{C}(\dot\gamma(-2nR_1))$. All trajectories from $x(y)$ to $z(w)$ cross $\partial\mathcal{C}(\dot\gamma(-2nR_1))$, so that we have the following estimates. For simplicity, we omit the domain $\mathcal{C}(\dot\gamma(-2(n+1)R_1-2))$ of the Green functions in the following paragraph.

$$\begin{aligned}
\frac{k_\lambda(x,y,z;n+1)}{k_\lambda(x,y,w;n+1)} - \underline{\theta}(n) &= \frac{G_\lambda(y,z)G_\lambda(x,w) - \underline{\theta}(n)G_\lambda(x,z)G_\lambda(y,w)}{G_\lambda(x,z)G_\lambda(y,w)} \\
= \quad & \frac{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} \left[G_\lambda(a,z)G_\lambda(b,w) - \underline{\theta}(n)G_\lambda(b,z)G_\lambda(a,w)\right]\, d\varpi_x^\lambda(b)d\varpi_y^\lambda(a)}{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} G_\lambda(a,w)G_\lambda(b,z)\, d\varpi_x^\lambda(b)d\varpi_y^\lambda(a)} \\
\sim_{(C_3C_0)^4} \quad & \frac{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} G_\lambda(y,a)G_\lambda(x,b)\left[G_\lambda(a,z)G_\lambda(b,w) - \underline{\theta}(n)G_\lambda(b,z)G_\lambda(a,w)\right] dadb}{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} G_\lambda(y,a)G_\lambda(x,b)G_\lambda(a,w)G_\lambda(b,z)\, dadb},
\end{aligned}$$

where we used Propositions 3.2 and 3.3 to write the last line. This is possible since both functions

$$G_\lambda(a,z)G_\lambda(b,w) - \underline{\theta}(n)G_\lambda(b,z)G_\lambda(a,w) \ \text{ and } \ G_\lambda(a,w)G_\lambda(b,z)$$

are positive harmonic in a and in b on a neighbourhood of size at least 1 of $\partial\mathcal{C}(\dot\gamma(-2nR_1))$. Using (3.5) with the point $x_n := \gamma(-(2n+1)R_1)$, we obtain

$$\begin{aligned}
& \frac{k_\lambda(x,y,z;n+1))}{k_\lambda(x,y,w;n+1)} - \underline{\theta}(n) \\
\sim_{(C_8C_3C_0)^4} \quad & \frac{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} G_\lambda(x_n,a)G_\lambda(x_n,b)\left[G_\lambda(a,z)G_\lambda(b,w) - \underline{\theta}(n)G_\lambda(b,z)G_\lambda(a,w)\right] dadb}{\int_{a,b\in\partial\mathcal{C}(\dot\gamma(-2nR_1))} G_\lambda(x_n,a)G_\lambda(x_n,b)G_\lambda(b,z)G_\lambda(a,w)\, dadb}.
\end{aligned}$$

Since the last line above doesn't depend on x and y, we have, setting $C' = (C_8C_3C_0)^8$,

$$\begin{aligned}
\overline{\theta}(n+1) - \underline{\theta}(n) &= \sup\{\frac{k_\lambda(x,y,z;(n+1))}{k_\lambda(x,y,w;(n+1))} - \underline{\theta}(n)\} \\
&\leq C' \inf\{\frac{k_\lambda(x,y,z;(n+1)))}{k_\lambda(x,y,w;(n+1))} - \underline{\theta}(n)\} \\
&= C'\left(\underline{\theta}(n+1) - \underline{\theta}(n)\right).
\end{aligned}$$

Applying an analogous argument to the function $\overline{\theta}(n) - \frac{k_\lambda(x,y,z;(n+1))}{k_\lambda(x,y,w;(n+1))}$, we get

$$\overline{\theta}(n) - \underline{\theta}(n+1) \leq C'\left(\overline{\theta}(n) - \overline{\theta}(n+1)\right).$$

Therefore, by adding the two inequalities and multiplying the results,

$$\overline{\theta}(n) - \underline{\theta}(n) \leq \left(\frac{C'-1}{C'+1}\right)^{n-1} \left(\overline{\theta}(1) - \underline{\theta}(1)\right) \leq C_8^2 \left(\frac{C'-1}{C'+1}\right)^{n-1}.$$

Since both $k(x,y,z)$ and $k(x,y,w)$ are 1 for $x=y$, $\underline{\theta} \leq 1 \leq \overline{\theta}$. Since the difference $\overline{\theta}(n) - \underline{\theta}(n)$ is small, they are both close to 1 and the ratio is between $\ln\underline{\theta}$ and $\ln\overline{\theta}$, which are of the same order as $\max\{\overline{\theta}-1, 1-\underline{\theta}\} \leq \overline{\theta} - \underline{\theta}$. Finally, we obtain constants C and $K<1$ such that, for all geodesic γ, all $\lambda \in [0,\lambda_0]$, all $x,y \in \mathcal{C}_{\pm1}(\dot\gamma(-2nR_1))$ and $z,w \in \mathcal{C}(\dot\gamma(2R_1))$

$$\left|\ln\frac{k_\lambda(x,y,z;n)}{k_\lambda(x,y,w;n)}\right| \leq CK^n. \tag{3.6}$$

□

Observe that the same proof also yields

Lemma 3.8. *There are constants $C > 1, K < 1$ such that for all geodesic γ, all $\lambda \in [0, \lambda_0]$, all $T > 1$, all $z, w \in \mathcal{C}(\dot\gamma(T))$, the hitting measure densities ϖ_z and ϖ_w satisfy, for all $a \in \partial\mathcal{C}(\dot\gamma(0))$,*

$$\ln \frac{\varpi_z}{\varpi_w}(a) \;\le\; CK^T.$$

Proof. The proof is the same as the proof of Lemma 3.7. The densities ϖ_y^k of the hitting measure on $\partial\mathcal{C}(\dot\gamma(2(N-n)R_1))$ starting from y, $y = z$ or w, satisfy:

$$\varpi_y^{n+1}(a_{n+1}) \;=\; \int_{\partial\mathcal{C}(\dot\gamma(2(N-n)R_1))} G_\lambda(a_n, a_{n+1})\, \varpi_y(a_n)\, da_n.$$

We can define as before

$$\overline{\theta}'_n \;:=\; \sup_{a\in\partial\mathcal{C}(\dot\gamma(2(N-n)R_1)} \frac{\varpi_z}{\varpi_w}(a); \quad \underline{\theta}'_n \;:=\; \inf_{a\in\partial\mathcal{C}(\dot\gamma(2(N-n)R_1)} \frac{\varpi_z}{\varpi_w}(a).$$

Replacing $G_\lambda(a_n, a_{n+1})$ by $G_\lambda(a_n, \gamma((2(N_n) - 1)R_1))G_\lambda(\gamma((2(N_n) - 1)R_1), a_{n+1})$, we obtain as above that

$$\overline{\theta}'_{n+1} - \underline{\theta}'_n \;\le\; C'(\underline{\theta}'_{n+1} - \underline{\theta}'_n).$$

The other inequality and the rest of the proof are the same as in Lemma 3.7. □

Proposition 3.9. *(1) Let $\xi \in \partial\widetilde{M}$ and $\lambda \le \lambda_0$. The following limit exists and defines a positive $(\Delta + \lambda)$-harmonic function*

$$k_\lambda(x, y, \xi) \;=\; \lim_{z\to\xi} k_\lambda(x, y, z),$$

which we call the Martin kernel.

(2) Fix $x, y \in \widetilde{M}$. There exist α and $C = C(\max\{d(x, y), 1\}) > 0$ such that for any $\lambda \in [0, \lambda_0]$,

$$\left|\ln \frac{k_\lambda(x, y, \xi)}{k_\lambda(x, y, \eta)}\right| \;\le\; C(d_x(\xi, \eta))^\alpha,$$

where d_x is the Gromov metric on $\partial\widetilde{M}$. Moreover, for $\beta < \alpha$, the function $\lambda \mapsto k_\lambda(x, y, \xi)$ is continuous from $[0, \lambda_0]$ into the space of β-Hölder continuous functions on $\partial\widetilde{M}$.

Proof. (1) It suffices to show it for a fixed $x = x_0$ and a sequence $z_n \to \xi$. Let γ be the geodesic going from x_0 to ξ. There is T such that $x_0, y \notin \mathcal{C}(\dot\gamma(T - 2R_1))$. As $n \to \infty$, $z_n \in \mathcal{C}(\dot\gamma(T_n + 2R_1))$, with $T_n \to \infty$. By Lemma 3.7, the sequence $k_\lambda(x_0, y, z_n)$ converges.

(2) Let γ be the geodesic such that $\gamma(0) = x, \gamma(+\infty) = \xi$. There is δ_0 depending only on the curvature bound such that if the Gromov distance $d_x(\xi, \eta)$ is smaller than δ_0, then there is $T = -C \ln d_x(\xi, \eta)$ such that ξ, η lie in the closure of $\mathcal{C}(\dot\gamma(T))$. We choose δ_0 small enough so that $T > \max\{d(x, y), 1\} + 4R_1$. Then, Lemma 3.7 applies to the limits $k_\lambda(x, y, \xi)$ and $k_\lambda(x, y, \eta)$ so that

$$\left|\ln \frac{k_\lambda(x, y, \xi)}{k_\lambda(x, y, \eta)}\right| \;\le\; C(d(x, y))K^{(-C \ln d_x(\xi,\eta))} \;=\; C(d_x(\xi, \eta))^\alpha,$$

where $\alpha = -C \ln K > 0$.

As λ varies, by Lemma 3.7, the functions $k_\lambda(x,y,z)$ are uniformly α-Hölder continuous on a neighborhood of ξ in $\widetilde{M} \cup \partial\widetilde{M}$. Moreover, $k_\lambda(x,y,z) \to k_{\lambda_0}(x,y,z)$ as $\lambda \to \lambda_0$. The continuity in λ follows for any $\beta < \alpha$. □

For $x,y,z \in \widetilde{M}$, denote

$$\theta_x^\lambda(y,z) \ := \frac{G_\lambda(y,z)}{G_\lambda(y,x)G_\lambda(x,z)}.$$

Proposition 3.10. *Fix $x \in \widetilde{M}, \xi \neq \eta \in \partial\widetilde{M}$. As $y \to \xi, z \to \eta$, the following limit exists and defines the Naïm kernel $\theta_x^\lambda(\xi,\eta)$:*

$$\theta_x^\lambda(\xi,\eta) \ := \lim_{y\to\xi, z\to\eta} \theta_x^\lambda(y,z) = \lim_{y\to\xi, z\to\eta} \frac{G_\lambda(y,z)}{G_\lambda(y,x)G_\lambda(x,z)}.$$

The limit is uniform in λ on the set of triples (x,ξ,η) with $d_x(\xi,\eta)$ bounded away from 0. For $v \in SM$, set $\theta_\lambda(v) := \theta^\lambda_{\gamma_{\widetilde{v}}(0)}(\widetilde{v}^-, \widetilde{v}^+)$. Then there is β such that the mapping $\lambda \mapsto \theta_\lambda$ is continuous from $[0,\lambda_0]$ to the space of β-Hölder continuous functions on SM.

Proof. Observe first that, by (3.5), for $d_x(y,z) = d(x,z)+d(x,y)-d(x,z)$ bounded away from 0, the functions $\theta_x^\lambda(y,z)$ are uniformly bounded. As before, by (3.7), the functions $y,z \mapsto \theta_x^\lambda(y,z)$ are uniformly α-Hölder continuous in y and in z as long as $d_x(y,z)$ remains bounded away from 0 and $\theta_x^\lambda(y,z) \to \theta_x^{\lambda_0}(y,z)$ as $\lambda \to \lambda_0$. The convergence and the continuity follow. Observe also that the function $\theta^\lambda_{\gamma_{\widetilde{v}}(0)}(\widetilde{v}^-, \widetilde{v}^+)$ is Γ-invariant and so θ_λ is indeed a function on SM. Since $d_{\gamma_{\widetilde{v}}(0)}(\widetilde{v}^-, \widetilde{v}^+) = 1$, the mapping $\lambda \mapsto \theta^\lambda_{\gamma_{\widetilde{v}}(0)}(\widetilde{v}^-, \widetilde{v}^+)$ is continuous from $[0,\lambda_0]$ to the space of β'-Hölder continuous functions on $\widetilde{SM}$ endowed with the metric coming from the identification with $\partial\widetilde{M} \times \partial\widetilde{M} \times \mathbb{R}$ for some $\beta' < \alpha$. This identification being itself Hölder continuous ([**AnS**] Proposition 2.1), the last statement of Proposition 3.10 follows.

□

For $v \in SM$, $x \in \widetilde{M}$, $\xi,\eta \in \partial\widetilde{M}$, we set

$$\theta(v) \ := \ \theta_{\lambda_0}(v), \quad \theta_x(\xi,\eta) := \theta_x^{\lambda_0}(\xi,\eta). \tag{3.7}$$

Fix $x,z \in \widetilde{M}, d(x,z) \geq 1$. The functions $x \mapsto k_\lambda(x,y,\xi)$ and $y \mapsto k_\lambda(x,y,z), \xi \in \partial\widetilde{M}$ are $\Delta + \lambda$ harmonic in y in a neighborhood of x. Let $v \in S_x\widetilde{M}$. The directional derivative $\partial_v k_\lambda(x,.,z)$ exists. Since $k(x,y,z)$ is a harmonic function of y away from z, by Proposition 2.7, $|\partial_v \ln k_\lambda(x,y,z)|_{y=x}| \leq \ln C_0$ where the constant $\ln C_0$ does not depend on $\lambda \in [0,\lambda_0]$. Following [**H1**] Lemma 3.2, we have:

Proposition 3.11. *For fixed $x_0 \in \widetilde{M}$ and $\widetilde{v} \in S_x\widetilde{M}$, the mapping $\xi \mapsto \partial_{\widetilde{v}} k_\lambda(x_0,y,\xi)|_{y=x_0}$ is α-Hölder continuous, uniformly in $\lambda \in [0,\lambda_0], \widetilde{v} \in S_x\widetilde{M}$. In particular, there is $\beta > 0$ such that the function $\lambda \mapsto \Phi_\lambda$ is continuous from $[0,\lambda_0]$ to the space of β-Hölder continuous functions on SM, where, for $v \in SM$, $\Phi_\lambda(v)$ is defined by*

$$\Phi_\lambda(v) := \partial_{\widetilde{v}} \ln k_\lambda(\gamma_{\widetilde{v}}(0),.,\gamma_{\widetilde{v}}(+\infty)) \ = \ \lim_{\varepsilon\to 0} \frac{1}{\varepsilon} \ln k_\lambda(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(\varepsilon), \gamma_{\widetilde{v}}(+\infty)),$$

where $\widetilde{v}$ is a lift of v.

Proof. Let $x_\varepsilon := exp_x(\varepsilon v)$ and take $z, d(x,z) \geq 2$. We have, for $v \in S_x\widetilde{M}$:

$$\partial_v k_\lambda(x,.,z) + \ln C_0 = \lim_{\varepsilon \to 0} \frac{\varepsilon^{-1}(G_\lambda(x_\varepsilon, z) - G_\lambda(x,z)) + \ln C_0 G_\lambda(x,z)}{G_\lambda(x,z)}.$$

Let γ be a geodesic with $\gamma(0) = x$. For $T > 2$ such that $z, w \in \mathcal{C}(\dot\gamma(T))$, and for $\varepsilon < 1$, we can write

$$\begin{aligned}\frac{G_\lambda(x_\varepsilon, z) - G_\lambda(x,z)}{\varepsilon} &+ \ln C_0 G_\lambda(x,z) \\ &= \int_{\partial\mathcal{C}(\dot\gamma(2))} \left[\frac{G_\lambda(x_\varepsilon, a) - G_\lambda(x,a)}{\varepsilon} + \ln C_0 G_\lambda(x,a)\right] \varpi_z(a)\, da\end{aligned}$$

and

$$\begin{aligned}\frac{G_\lambda(x_\varepsilon, w) - G_\lambda(x,w)}{\varepsilon} &+ \ln C_0 G_\lambda(x,w) \\ &\int_{\partial\mathcal{C}(\dot\gamma(2))} \left[\frac{G_\lambda(x_\varepsilon, a) - G_\lambda(x,a)}{\varepsilon} + \ln C_0 G_\lambda(x,a)\right] \varpi_w(a)\, da,\end{aligned}$$

where we wrote the hitting measure on $\partial\mathcal{C}(\dot\gamma(2))$ starting from $y = z$ or w as $\varpi_y(a)\, da$. As $\varepsilon \to 0$ the expression between brackets is positive, converge towards $\partial_v G_\lambda(x,a) + \ln C_0 G_\lambda(x,a)$ and is dominated by $2 \ln C_0 G_\lambda(x,a)$, which is integrable with respect to $\varpi_z(a) da$ and $\varpi_w(a) da$. Since $z, w \in \mathcal{C}(\dot\gamma(T))$ we have, by Lemma 3.8, that for any $a \in \partial\mathcal{C}(\gamma(2))$,

$$\ln \frac{\varpi_z}{\varpi_w}(a) \ \leq \ CK^T.$$

By choosing the best possible T such that $z, w \in \mathcal{C}(\dot\gamma(T))$, we obtain that for any $a \in \partial\mathcal{C}(\gamma(2))$,

$$\frac{\varpi_z}{\varpi_w}(a) \sim^{C(d_x(z,w))^\alpha} 1, \tag{3.8}$$

if $d_x(z,w)$ is small enough. The estimate (3.8) is preserved in the integral and in the limit as $\varepsilon \to 0$ so that

$$\frac{\partial_v k_\lambda(x,.,z) + \ln C_0}{\partial_v k_\lambda(x,.,w) + \ln C_0} \sim^{C(d_x(z,w))^\alpha} 1,$$

if $d_x(z,w)$ is small enough. This proves that the mapping $\xi \mapsto \partial_v k_\lambda(x,.,\xi)$ is α-Hölder continuous, uniformly in $\lambda \in [0, \lambda_0], v \in S_x\widetilde{M}$. The other statements of the Proposition follow.

□

Corollary 3.12. *Set $P(\lambda_0)$ for the pressure of the function $\varphi_{\lambda_0} := -2\Phi_{\lambda_0}$. Then, $P(\lambda_0) \leq 0$.*

Indeed we know by Corollary 2.11 that the pressure of the function φ_λ is negative, and by Proposition 3.11 that the mapping $\lambda \mapsto \varphi_\lambda$ is continuous at λ_0.

Corollary 3.13. *The measures μ_λ and the normalising constants $\Omega_\lambda, \Upsilon_\lambda$ are continuous functions of λ as $\lambda \to \lambda_0$ in $[0, \lambda_0]$.*

Proof. Indeed, the measures μ_{λ_0} satisfies the conditions in Proposition 2.6 and Ω_{λ_0} satisfies the expression (2.9). Since all the functions involved are continuous by Proposition 3.9 and Proposition 3.11, Corollary follows. The argument is the same for Υ_λ. □

We can now prove Theorem 1.3 giving the exponential decay of $G_{\lambda_0}(x, y)$ with the distance. More precisely, we have:

Proposition 3.14. *Let $\tau_0 := \inf\{\int \Phi_{\lambda_0}\, dm\}$, where the infimum is taken over all g-invariant probability measures. Then, $\tau_0 > 0$ and*

$$\lim_{R\to\infty} -\frac{1}{R} \ln \max\{G_{\lambda_0}(x, y) : d(x, y) = R\} \ = \tau_0.$$

Proof. First we prove that $\tau_0 > 0$. Suppose m_1 attains $\sup \int \varphi_{\lambda_0} dm$. Suppose $\int \varphi_{\lambda_0}\, dm_1 \geq 0$. Then $h_{m_1} + \int \varphi_{\lambda_0}\, dm_1 \geq 0$. However, we know that $P(\varphi_{\lambda_0}) \leq 0$, thus $h_{m_1} = 0$ and $\int \varphi_{\lambda_0} dm_1 = 0$, and therefore m_1 is the equilibrium state of φ_{λ_0}. This is a contradiction since $h_{m_1} > 0$ if m_1 is the equilibrium state of the Hölder continuous function φ_{λ_0}. This proves that $\sup\{\int \varphi_{\lambda_0}\, dm\} < 0$ and thus that τ_0 is positive.

Let $0 < \tau' < \tau_0$. We have

$$\lim_{t\to\infty} \frac{1}{t} P(-t\Phi_{\lambda_0}) \ = \ -\tau_0,$$

so we can find T large enough that $-P(-T\Phi_{\lambda_0}) > T\tau'$. By Corollary 2.14, we get that there exists a constant $C(T)$ such that for all $R \geq 1, x \in \widetilde{M}$,

$$e^{-RP(-T\Phi_{\lambda_0})} \int_{S(x,R)} G^T_{\lambda_0}(x, z) dz \ \leq C(T).$$

Set $\tau(R) := \max\{-\frac{1}{R} \ln G_{\lambda_0}(x, z) : d(x, z) = R\}$. By compactness, there exist x, y with $d(x, y) = R$ and $G_{\lambda_0}(x, y) = e^{-R\tau(R)}$. We have, for $z \in S(x, R), d(y, z) \leq 1$,

$$G_{\lambda_0}(x, z) \geq C_0^{-1} e^{-R\tau(R)} \quad \text{ and thus } \quad G^T_{\lambda_0}(x, z) \geq C_0^{-T} e^{-TR\tau(R)}.$$

Therefore, we have for all $R \geq 1$,

$$C(T) \geq e^{RT\tau'} \int_{S(x,R)\cap B(y,1)} G^T_{\lambda_0}(x, z) dz \ \geq C_0^{-T} e^{RT(\tau' - \tau(R))}.$$

This is possible only if $\liminf_R \tau(R) \geq \tau'$. Since $\tau' < \tau_0$ was arbitrary, this proves that

$$\liminf_{R\to\infty} -\frac{1}{R} \ln \max\{G_{\lambda_0}(x, y) : d(x, y) = R\} \ \geq \tau_0.$$

Conversely, for all $\varepsilon > 0$, there exists a closed geodesic of length ℓ, say, such that for v tangent to that geodesic,

$$\int_0^\ell \Phi_{\lambda_o}(g_s v)\, ds \ \leq \ (\tau_0 + \varepsilon)\ell.$$

Let $\widetilde{v}$ be a lift of v. The geodesic $\gamma_{\widetilde{v}}$ is a periodic axis and for all $j \in \mathbb{N}$,

$$k_{\lambda_0}(\gamma_{\widetilde{v}}(j\ell), \gamma_{\widetilde{v}}((j+1)\ell), \gamma_{\widetilde{v}}(+\infty)) \ \leq \ e^{(\tau_0+\varepsilon)\ell}.$$

By Lemma 3.7, we have:

$$\frac{G_{\lambda_0}(\gamma_{\widetilde{v}}(j\ell), \gamma_{\widetilde{v}}(N\ell))}{G_{\lambda_0}(\gamma_{\widetilde{v}}((j+1)\ell), \gamma_{\widetilde{v}}(N\ell))} \ \geq \ \left(1 + CK^{(N-j)\ell}\right)^{(-1)} e^{-(\tau_0+\varepsilon)\ell}$$

and therefore

$$\frac{G_{\lambda_0}(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(N\ell))}{G_{\lambda_0}(\gamma_{\widetilde{v}}((0)\ell), \gamma_{\widetilde{v}}(\ell))} = \prod_{j=0}^{N-2} \frac{G_{\lambda_0}(\gamma_{\widetilde{v}}(j\ell), \gamma_{\widetilde{v}}(N\ell))}{G_{\lambda_0}(\gamma_{\widetilde{v}}((j+1)\ell), \gamma_{\widetilde{v}}(N\ell))} \ \geq Ce^{-N(\tau_0+\varepsilon)\ell}.$$

This shows that, for all $\varepsilon > 0$,

$$\limsup_{R\to\infty} -\frac{1}{R} \ln \max\{G_{\lambda_0}(x,y) : d(x,y) = R\} \ \leq \tau_0 + \varepsilon.$$

□

Lemma 3.15. *There exists $C > 0$ such that for any $\lambda \in [0, \lambda_0]$ and x, ξ, η, there exists $x_0 \in [\xi, \eta]$ such that if y is in the geodesic ray from x to ξ, then*

$$\frac{k_\lambda(x,y,\eta)}{k_\lambda(x,y,\xi)} \leq C \min(1, G_\lambda(y,x_0)^2).$$

Proof. We first claim that by δ-hyperbolicity, there exist points $x_0 \in [\xi,\eta], x_1 \in [x,\eta], x_2 \in [y,\eta], x_3 \in [x,\xi]$ such that the distance between them is bounded above by 3δ. Indeed, for x' in the geodesic from η to ξ, the distance function $x' \mapsto d(x',[x,\xi])$ is a decreasing function. Let x' the first point where $d(x',[x,\xi]) \leq \delta$ and choose $x_0 \in [x',\eta]$ to be the point δ-apart from x'. By definition, $\delta < d(x_0,[x,\xi]) \leq 2\delta$, thus there exists $x_3 \in [x,\xi]$ of distance 2δ-close to x_0. Choose $x_1 \in [x,\eta], x_2 \in [y,\eta]$ δ-close to x_0. The claim follows.

Now choose θ_0 such that if $\angle_x(\xi,\eta) \leq \theta_0$, then x is R_0-apart from $x_0, \cdots, x_3$. By Theorem 3.4 (which gives estimates up to C_1) and Harnack inequality (which gives estimates up to C_H) , the following estimates follow.

Let $[x,\xi] \ni w \to \xi$ and $[x,\eta] \ni z \to \eta$. Let us write $G(x,y) = G_\lambda(x,y)$ for simplicity.

(1) If $d(x,y) < d(x,x_3) - R_0 - 3\delta$, then

$$\frac{k_\lambda(x,y,z)}{k_\lambda(x,y,w)} = \frac{G(y,z)G(x,w)}{G(x,z)G(y,w)} \sim^{(C_1C_H)^4} \frac{G(y,x_0)G(x_0,z)}{G(x,x_0)G(x_0,z)} \frac{G(x,x_0)G(x_0,w)}{G(y,x_0)G(x_0,w)} = 1.$$

(2) If $d(x,y) > d(x,x_3) + R_0 + 3\delta$, then

$$\frac{k_\lambda(x,y,z)}{k_\lambda(x,y,w)} \sim^{(C_1C_H)^4} \frac{G(y,x_0)G(x_0,z)}{G(x,x_0)G(x_0,z)} \frac{G(x,x_0)G(x_0,y)G(y,w)}{G(y,w)} = G(x_0,y)^2.$$

(3) If $|d(x,y) - d(x,x_3)| < R_0 + 3\delta$, then

$$\frac{k_\lambda(x,y,z)}{k_\lambda(x,y,w)} = \frac{G(y,z)G(x,w)}{G(x,z)G(y,w)} \sim^{C_1C_H^2} \frac{G(y,z)}{G(x,y)G(y,z)} \frac{G(x,y)G(y,w)}{G(y,w)} = 1.$$

For x, ξ, η such that $\angle_x(\xi, \eta) > \theta_0$, if $d(y, x) > R_0$ and $d(y, x_2) > R_0$, then

$$\frac{k_\lambda(x,y,z)}{k_\lambda(x,y,w)} = \frac{G(y,z)G(x,w)}{G(x,z)G(y,w)} \sim^{C_1^2} \frac{G(y,x_2)G(x_2,z)}{G(x,z)}\frac{G(x,y)G(y,w)}{G(y,w)} \sim^{C_H^2} G^2(x_0,y).$$

Otherwise, $d(x, x_2)$ and thus $d(y, x), d(y, x_2)$ are all bounded by a constant depending on θ_0, thus the Lemma follows from Proposition 2.7. □

Remark 3.16. Note that from the proof, if the angle θ between η and ξ is bounded away from zero, then $d(x, x_0)$ is bounded above.

Now let us prove Theorem 1.4. Recall that Martin compactification of the operator $\Delta + \lambda_0$ is given by all possible limits of $k_{\lambda_0}(x, y, z)$ as $z \to \infty$. Proposition 3.9 and its proof show that there is a continuous mapping from the geometric compactification of $\widetilde{M}$ onto the Martin compactification. So it suffices to show that this mapping is one-to-one. If $\eta \neq \xi$, by Lemma 3.15, $k_{\lambda_0}(x, y, \eta)/k_{\lambda_0}(x, y, \xi) \to 0$ as $y \to \xi$ and thus $k_{\lambda_0}(x, ., \xi)$ does not coincide with $k_{\lambda_0}(x, ., \eta)$.

The decomposition of positive $(\Delta + \lambda_0)$-harmonic functions follows then by general Martin theory.

Since by Proposition 3.14, $G_{\lambda_0}(x_0, \cdot)$ goes to 0 at infinity uniformly, we get the following estimate for small $d(x, y)$:

Corollary 3.17. *There is a constant* C *such that for* $0 < d(x, y) \leq 1$,
(3.9)

$$C^{-1} \leq (d(x,y))^{m-2} G_{\lambda_0}(x,y) \leq C \text{ for } m > 2, \quad C^{-1} \leq \frac{G_{\lambda_0}(x,y)}{|\ln d(x,y)|} \leq C \text{ for } m = 2.$$

Proof. Observe that, for $x \neq y$,

$$G_{\lambda_0}(x,y) = \int_0^1 e^{\lambda_0 t} \wp(t,x,y)\,dt + \int_1^\infty e^{\lambda_0 t} \wp(t,x,y)\,dt$$

and that the last term is uniformly bounded for $d(x, y) \leq 1$:

$$\begin{aligned} \int_1^\infty e^{\lambda_0 t} \wp(t,x,y)\,dt &= e^{\lambda_0} \int_{\widetilde{M}} \wp(1,x,z) G_{\lambda_0}(z,y)\,d\mathrm{Vol}(z) \\ = e^{\lambda_0} \int_{B(x,2)} \wp(1,x,z) G_{\lambda_0}(z,y)\,d\mathrm{Vol}(z) &+ e^{\lambda_0} \int_{\widetilde{M} \setminus B(x,2)} \wp(1,x,z) G_{\lambda_0}(z,y)\,d\mathrm{Vol}(z) \\ \leq e^{\lambda_0} \max_{B(x,2)} \wp(1,x,z) \int_{B(y,3)} G_{\lambda_0}(z,y)\,d\mathrm{Vol}(z) &+ e^{\lambda_0} \max_{d(z,y)\geq 1} G_{\lambda_0}(z,y). \end{aligned}$$

We used Lemma 2.1 to bound uniformly $\int_{B(y,3)} G_{\lambda_0}(z, y)\,d\mathrm{Vol}(z)$ and Proposition 3.14 to show that $\max_{d(z,y)\geq 1} G_{\lambda_0}(z, y) < \infty$.
Since the curvature is bounded, it follows from [**Mv**] that for $0 < t \leq 1, 0 < d(x, y) \leq 1$

$$\wp(t,x,y)(4\pi t)^{m/2} e^{-\frac{d(x,y)^2}{4t}} \sim^C 1.$$

Corollary 3.17 follows by integration in t. □

Corollary 3.18. *There is a constant* C *such that for* $0 < d(x,y) \le 1$,

$$\int_{B(x_k,4)} G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y)\,d\mathrm{Vol}(z) \;\le\; CG_{\lambda_0}(x_k,y)$$

Indeed, by Corollary 3.17, it suffices to show that there is a constant C such that

$$\begin{aligned}\int_{B(x,4)} \frac{d\mathrm{Vol}(z)}{(d(x,z)d(y,z))^{m-2}} &\le \frac{C}{d(x,y)^{m-2}} \qquad \text{for } m>2,\\ \int_{B(x,4)} |\ln d(x,z)\ln(dy,z)|d\mathrm{Vol}(z) &\le C|\ln d(x,y)| \qquad \text{for } m=2.\end{aligned}$$

The statement reduces to the Euclidean case, where it can be shown by direct computation.

4. Renewal theory

In this section, we use uniform mixing of the geodesic flow g_t we obtained from Section 7 to control the convergence in Proposition 2.10 as λ goes to λ_0. Throughout the section, let us denote $\chi(t) := 1$ for $|t| \le 1/2$ and 0 otherwise. Let $\chi_{\delta'}(t) = \chi(t/\delta')$. Let $\psi(t) := \max\{1-|t|, 0\}$.

Thanks to Proposition 3.11, for λ close to λ_0, the functions φ_λ are close to φ_{λ_0} in the space $\mathcal{K}_\alpha$ of α-Hölder continuous functions, for some $\alpha > 0$ (see Section 7.1 for definition of $\mathcal{K}_\alpha$). We can therefore apply Proposition 7.3 and Corollary 7.4 to the equilibrium measure m_λ associated to φ_λ to obtain the following propositions.

Proposition 4.1. *There exist* $\alpha > 0$ *and* $\delta_0 > 0$ *with the following property. For every* $\varepsilon > 0$, $f, h \in \mathcal{K}_\alpha$ *positive* α*-Hölder continuous functions, there exists* $t_0 = t_0(f,h,\varepsilon)$, *such that for* $t \ge t_0$, *for any* $\lambda \in [\lambda_0-\delta_0, \lambda_0]$,

$$\int_{SM} fh\circ g_t\,dm_\lambda \sim^{1+\varepsilon} \int_{SM} f\,dm_\lambda \int_{SM} h\,dm_\lambda.$$

Indeed, t_0 *depends only on* $(\varepsilon, ||f||_\alpha, ||h||_\alpha, \inf_\lambda \int f\,dm_\lambda, \inf_\lambda \int h\,dm_\lambda)$, *in particular is independent of* $\lambda \in [\lambda_0-\delta_0, \lambda_0]$.

Proposition 4.2. *There exist* $\alpha' > 0$ *and* $\delta_0' > 0$ *with the following property. For every* $\varepsilon > 0$, $f, u, h \in \mathcal{K}_{\alpha'}$ *positive* α'*-Hölder continuous functions, there exists* $t_0' = t_0'(f,u,h,\varepsilon)$, *such that for* $t \ge t_0'$, *for any* $\lambda \in [\lambda_0-\delta_1, \lambda_0]$,

$$\frac{1}{t}\int_0^t \left[\int f\cdot(u\circ g_s)\cdot(h\circ g_t)\,dm_\lambda\right] ds \sim^{1+\varepsilon} \int f\,dm_\lambda \int u\,dm_\lambda \int h\,dm_\lambda.$$

Indeed, t_0' *depends only on* $(\varepsilon, ||f||_{\alpha'}, \|u\|_{\alpha'}, ||h||_{\alpha'}, \inf_\lambda \int f\,dm_\lambda, \inf_\lambda \int u\,dm_\lambda, \inf_\lambda \int h\,dm_\lambda)$, *in particular is independent of* $\lambda \in [\lambda_0-\delta_0', \lambda_0]$.

4.1. **Integral on large spheres with respect to Green functions.** Let us introduce some more notations: for $x \neq z \in \widetilde{M}$, denote by v_z^x the unit vector in $S_z\widetilde{M}$ pointing towards x. The mapping $z \mapsto v_z^x$ identifies $\widetilde{M} \setminus \{x\}$ with a subset of $S\widetilde{M}$.

Theorem 4.3. *Given $\varepsilon' > 0$ and positive Hölder continuous functions f, h on SM, there exist $R(f,h,\varepsilon')$ and $\delta(f,h,\varepsilon')$ such that if $R > R(f,h,\varepsilon')$ and $\lambda \in [\lambda_0 - \delta(f,h,\varepsilon'), \lambda_0]$, for all $x \in \widetilde{M}$,*

$$(4.1) \qquad e^{-RP(\lambda)} \int_{S(x,R)} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x,y) dy \qquad \sim^{(1+\varepsilon')^3}$$

$$\Omega_\lambda \int_{\partial\widetilde{M}} f(p \circ \pi_x^{-1}\xi) d\mu_x^\lambda(\xi) \int_{M_0} \left(\int_{\partial\widetilde{M}} h(p \circ \pi_y^{-1}\xi) d\mu_y^\lambda(\xi) \right) d\,Vol(y).$$

The rest of Section 4.1 is devoted to the proof of Theorem 4.3.

Fix f, h positive and Hölder continuous. We choose $\delta_0' > 0$ such that, if $R > 1$ and $|R - R'| < \delta_0'$, then, for all $x \in \widetilde{M}$ and $\lambda \in [\lambda_0 - \delta(f,h,\varepsilon'), \lambda_0]$,

(4.2)

$$e^{-RP(\lambda)} \int_{S(x,R)} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x,y) dy \sim^{1+\varepsilon'} e^{-R'P(\lambda)} \int_{S(x,R')} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x,y) dy.$$

Then

$$\delta'\,(4.1) \ \sim^{(1+\varepsilon')^2} \int_{\mathbb{R}} \chi_{\delta'}(s-R) e^{-sP(\lambda)} \left(\int_{S(x,s)} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x,y) dy \right) ds.$$

$$= \int_{\widetilde{M}} \chi_{\delta'}(d(x,y) - R) e^{-d(x,y)P(\lambda)} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x,y) d\mathrm{Vol}(y)$$

$$= \int_{M_0} \left(\sum_{\{(v,t): v \in S_{px}M \cap g_{-t}S_{py}M\}} \chi_{\delta'}(t-R) f(v) h(-g_t v) e^{-tP(\lambda)} G_\lambda^2(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t)) \right) d\mathrm{Vol}(y).$$

Recall from Section 2 that

$$\frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) = e^{-tP(\lambda)} k_\lambda^2(\gamma_{\widetilde{v}}(t), \gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(\infty)).$$

By Proposition 3.10, for given ε, if R is large enough (depending on ε) and $|t - R| \leq \delta' \leq 1$,

$$\begin{aligned} \frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) \quad &= \quad e^{-tP(\lambda)} \lim_{z \to \widetilde{v}^+} \frac{G_\lambda^2(\gamma_{\widetilde{v}}(0), z)}{G_\lambda^2(\gamma_{\widetilde{v}}(t), z) G_\lambda^2(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t))} G_\lambda^2(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t)) \\ &\sim^{1+\varepsilon} \quad e^{-tP(\lambda)} \theta_\lambda^2(g_t v)\, G_\lambda^2(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t)), \end{aligned}$$

where the approximation is uniform in $g_t v$ and λ. Therefore, when replacing

$$e^{-tP(\lambda)} G_\lambda^2(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t)) \quad \text{by} \quad \frac{1}{\theta_\lambda^2(g_t v)} \frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v),$$

for R large enough and $|t-R| \leq \delta'$, uniformly in x, y and λ, we loose a factor $(1+\varepsilon')^{\pm 1}$. Remark that $\theta_\lambda(v) = \theta_\lambda(-v)$.

We are reduced to show:

Proposition 4.4. *Given $\varepsilon' > 0$ and positive Hölder continuous functions f, h on SM, there exist $R_0 = R_0(f, h, \varepsilon')$, $\delta = \delta(f, h, \varepsilon') > 0$ and $\delta', 0 < \delta' < \delta'_0$, such that for $R \geq R_0$, all $x, y \in \widetilde{M}$ and all $\lambda \in [\lambda_0 - \delta, \lambda_0]$,*

$$\sum_{\{(v,t): v \in S_{px}M \cap g_{-t}S_{py}M\}} \chi_{\delta'}(R-t) f(v) (\theta^{-2}h)(-g_t v) \frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v)$$

$$\sim^{1+\varepsilon'} \ \Omega_\lambda \delta' \left(\int_{S_{px}M} f(v)\, d\mu_\lambda^{uu}(v) \right) \left(\int_{S_{py}M} h(-u)\, d\mu_\lambda^{ss}(u) \right).$$

The last line in Proposition 4.4 is the same as

$$\delta' \Omega_\lambda \int_{\partial \widetilde{M}} f(p \circ \pi_x^{-1} \xi) d\mu_x^\lambda(\xi) \int_{\partial \widetilde{M}} h(p \circ \pi_y^{-1} \xi) d\mu_y^\lambda(\xi)$$

by (2.12) and (2.13).

Theorem 4.3 follows from Proposition 4.4 and the previous discussion by integrating the approximation in y over a fundamental domain M_0.

Proof. We combine ideas of [**M1**] and Section III in [**L1**]. Choose ε such that $(1+\varepsilon)^{62} \leq 1 + \varepsilon'$. Proposition follows from Proposition 4.1 applied to the non-negative Hölder continuous functions $F_\lambda^\pm, H_\lambda^\pm$ with the property that there exist constants $C, \alpha, \beta_0, \beta'_0, \gamma$ such that for all $x, y \in \widetilde{M}$ and all $\lambda \in [0, \lambda_0]$, the following (1)-(5) holds.

(1) $||F_\lambda^\pm||_\alpha < C, ||H_\lambda^\pm||_\alpha < C,$
(2) $\int F_\lambda^\pm dm_\lambda > C^{-1}, \int H_\lambda^\pm dm_\lambda > C^{-1}.$
(3)

$$\begin{aligned} \Omega_\lambda \delta' \beta_0 (1+\varepsilon)^{-14} \int_{S_{px}M} f(v) d\mu_\lambda^{uu}(v) \ &\leq \ \int F_\lambda^- dm_\lambda \\ \leq \ \int F_\lambda^+ dm_\lambda \ &\leq \ \Omega_\lambda \delta' \beta_0 (1+\varepsilon)^{14} \int_{S_{px}M} f(v) d\mu_\lambda^{uu}(v). \end{aligned}$$

(4)

$$\begin{aligned} \Omega_\lambda \gamma \beta'_0 (1+\varepsilon)^{-14} \int_{S_{py}M} h(-u) d\mu_\lambda^{ss}(u) \ &\leq \ \int H_\lambda^- dm_\lambda \\ \leq \ \int H_\lambda^+ dm_\lambda \ &\leq \ \Omega_\lambda \gamma \beta'_0 (1+\varepsilon)^{14} \int_{S_{py}M} h(-u) d\mu_\lambda^{ss}(u). \end{aligned}$$

(5) There is $R(\varepsilon)$ such that for $R \geq R(\varepsilon)$,

$$\begin{aligned} & (1+\varepsilon)^{-32} \int F_\lambda^- H_\lambda^- \circ g_R dm_\lambda \\ \leq \quad & \Omega_\lambda \gamma \beta_0 \beta_0' \sum_{\{(v,t): v \in S_{px}M \cap g_{-t} S_{py} M\}} \chi_{\delta'}(R-t) f(v) (\theta^{-2} h)(-g_t v) \frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) \\ \leq \quad & (1+\varepsilon)^{32} \int F_\lambda^+ H_\lambda^+ \circ g_R dm_\lambda. \end{aligned}$$

We choose $\delta' < \delta_0'$ and such that, for all $\xi \in \partial \widetilde{M}$, all $\lambda \in [0, \lambda_0]$, all

$$\frac{\theta^2(v')}{\theta^2(v)}, k_\lambda(x, x', \xi), e^{-P\beta(x,x',\xi)} \sim^{1+\varepsilon} 1 \text{ for } d(v,v') < 2\delta', d(x,x') < 2\delta',$$

where $P := \inf_{\lambda \in [0,\lambda_0]} P(\lambda) < 0$.

The functions $F_\lambda^\pm(v), H_\lambda^\pm(u)$ will approximate $\theta_\lambda^{-2} f(v), \theta_\lambda^{-2} h(-u)$ respectively, on the δ-neighborhoods $N_\delta(S_{px}M)$, $N_\delta(S_{py}M)$ of $S_{px}M$, $S_{py}M$, respectively.

For $w \in N_{\delta'}(S_{px}M)$, there exist a unique $v \in S_{px}M$, and $v' \in W_{loc}^{ss}(v), t$ such that $v' = \gamma_t w$. Similarly, if $w \in N_\gamma(S_{py}M)$, then there exists a unique triple $(u, u', s), u \in S_{py}M, u' \in W_{loc}^{uu}(u)$ such that $u' = g_s(w)$.

By the Hölder regularity of the strong stable and the strong unstable foliations, the systems of coordinates (v, v', t) (respectively (u, u', t)) are Hölder continuous, uniformly in x and y.

Step 1. There exist $\beta_0, \beta_0' > 0$ and non-negative Hölder continuous functions $a_\pm, b_\pm$ supported on $N_\delta S_{px}M$, $N_\delta S_{py}M$, respectively, such that for all $v \in S_{px}M$ and $u \in S_{py}M$,

$$\int_{W_{loc}^{ss}(v)} a_\pm(w)\, d\mu_\lambda^{ss}(w) = \beta_0 (1+\varepsilon)^{\pm 1}, \quad \int_{W_{loc}^{uu}(u)} b_\pm(w)\, d\mu_\lambda^{uu}(w) = \beta_0'(1+\varepsilon)^{\pm 1}. \tag{4.3}$$

Moreover, the Hölder exponent and the Hölder coefficient of $a_\pm, b_\pm$ are bounded uniformly in x, y, λ.

Lemma 4.5. *Let*

$$h_{r,v,\lambda} = \int_{W_{loc}^{ss}(v)} \psi\left(\frac{d_{ss}(v, v')}{r} \right) d\mu_\lambda^{ss}(v').$$

The map $(r, v, \lambda) \mapsto h_{r,v,\lambda}$ is continuous in r, v and λ. For a fixed r, the function $v \mapsto h_{r,v,\lambda}$ is Hölder continuous, uniformly in $\lambda \in [0, \lambda_0]$. As r varies from 0 to δ', the function $r \mapsto h_{r,v,\lambda}$ is increasing and admits right and left derivatives that are are bounded below by a positive constant uniformly in v, λ and r away from zero.

Proof. The continuity is as in Margulis's Lemma 7.1 in [**M2**](p. 51). The proof also yields Hölder continuity in v. Indeed, $W_{loc}^{ss}(v)$ depends on v in a Hölder continuous way and if v_1, v_2 are close, the holonomy H_1^2 from $W_{loc}^{ss}(v_1)$ to $W_{loc}^{ss}(v_2)$ is Hölder continuous, and satisfies for $v_1', v_1'' \in W_{loc}^{ss}(v_1)$,

$$d(v_2, H_1^2 v_1) \leq C(d(v_1, v_2))^\alpha, \text{ and } |d(H_1^2 v_1', H_1^2 v_1'') - d(v_1', v_1'')| \leq C(d(v_1', v_1''))^\alpha.$$

Moreover the logarithm of the Radon Nikodym derivatives of the measure $(H_1^2)_*\mu_\lambda^{ss}(v_2')$ with respect to $\mu_\lambda^{ss}(v_2')$ is also, uniformly in λ, proportional to $(d(v_2, H_1^2 v_1))^\alpha$ (Proposition 3.9). Since $d(v_2, H_1^2 v_1) \leq C(d(v_1, v_2))^\alpha$, we can report in the definition of $h_{r,v,\lambda}$ and see that, for v_1, v_2 close,

$$|h_{r,v_1,\lambda} - h_{r,v_2,\lambda}| \ \leq \ C(r)(d(v_1, v_2))^\alpha,$$

where the constant $C(r)$ is uniform in $\lambda \in [0, \lambda_0]$ and goes to infinity as $r \to 0$.

Direct computation shows that, as r varies from 0 to δ', the function $r \mapsto h_{r,v,\lambda}$ is increasing and admits left and right derivatives given by

$$\frac{\partial}{\partial r} h_{r,v,\lambda}|_{r-} = \lim_{r'<r, r'\to r} \int_{W_{loc}^{ss}(v)} \frac{1}{r'} d_{ss}(v,v') \chi_{d(v,\cdot)\leq r'}(v')\ d\mu_\lambda^{ss}(v')$$

and

$$\frac{\partial}{\partial r} h_{r,v,\lambda}|_{r+} = \int_{W_{loc}^{ss}(v)} \frac{1}{r} d_{ss}(v,v') \chi_{d(v,\cdot)\leq r}(v')\ d\mu_\lambda^{ss}(v').$$

In particular, the derivatives are bounded from below by a positive constant uniformly in v, λ and r away from 0. $\square$

For given β, choose $r_\lambda^\pm(v,\beta)$ such that $h_{r_\lambda^\pm(v,\beta),v,\lambda} = \beta(1+\varepsilon)^{\pm 1}$. Now choose β_0 so that $r_\lambda^\pm(v,\beta_0) < \varepsilon\delta'/2$ for all v and λ. Set $r_\lambda^\pm(v) := r_\lambda^\pm(v,\beta_0)$. By the Implicit function theorem with Hölder coefficients, the functions $r_\lambda^\pm(v)$ are Hölder continuous uniformly in λ for $\lambda \in [\lambda_0 - \delta(\varepsilon), \lambda_0]$ and v.

Now for $w = (v, v', t) \in N_\delta(S_{px}M), \lambda \in [\lambda_0 - \delta(\varepsilon), \lambda_0]$, define

$$a_\lambda^\pm(w) = \psi\left(\frac{d_{ss}(v,v')}{r_\lambda^\pm(v)}\right).$$

Properties similar to Lemma 4.5 holds for the function

$$(r, u, \lambda) \mapsto h_{r,u,\lambda} = \int_{W_{loc}^{uu}(u)} \psi\left(\frac{d_{uu}(u,u')}{r}\right) d\mu_\lambda^{uu}(u'),$$

thus we can define $r_\lambda^\pm(u)$ analogously: β_0' is chosen so that $r_\lambda^\pm(u, \beta_0') < \varepsilon\delta'/2$ and $r_\lambda^\pm(u)$ is such that $h_{r_\lambda^\pm(u),u,\lambda} = \beta_0'(1+\varepsilon)^{\pm 1}$. For $w = (u, u', s) \in N_\gamma(S_{py}M)$, define

$$b_\lambda^\pm(w) = \psi\left(\frac{d_{uu}(u,u')}{r_\lambda^\pm(u)}\right).$$

The functions $a^\pm, b^\pm$ satisfy the properties of Step 1. $\square$

Step 2. Definition of $F_\lambda^\pm, H_\lambda^\pm$ and Property (1)

Consider Lipschitz continuous $\chi_\pm(t)$ on $\mathbb{R}$ such that, for all $t \in \mathbb{R}$,

$$\chi_{(1+\varepsilon)^{-2}}(t) \leq \chi_-(t) \leq \chi_{(1+\varepsilon)^{-1}}(t) \leq \chi(t) \leq \chi_{(1+\varepsilon)}(t) \leq \chi_+(t) \leq \chi_{(1+\varepsilon)^2}(t).$$

Now for $w = (v, v', t)$, define

$$F_\lambda^\pm(w) = \chi_\pm(t/\delta') a_\pm(v')(\theta_\lambda^{-2} f)(v)$$

and for $w = (u, u', s)$,

$$H_\lambda^\pm(w) = \chi_\pm(s/\gamma) b_\pm(u')(\theta_\lambda^{-2} h)(-u),$$

for some $\gamma < \delta'\varepsilon/2$.

Recall that the systems of coordinates (v, v', t) and (u, u', s) are Hölder continuous uniformly in x and y. The functions $F_\lambda^\pm, H_\lambda^\pm$ in those coordinates are compositions of Hölder continuous functions (ψ, f, h) and of the functions $r_\pm$ that depend on v in a Hölder continuous way, uniformly in $\lambda \in [0, \lambda_0]$ by Step 1, which proves Property (1).

Step 3. Properties (2), (3) and (4)

Recall that under Hopf parametrisation introduced in Section 2, if we let $x_0 = x$, the lift $\widetilde{m}_\lambda$ of m_λ to $\widetilde{SM}$ is given by

$$d\widetilde{m}_\lambda(\xi, \eta, t) = \Omega_\lambda \theta_x^2(\xi, \eta) e^{2P(\lambda)(\xi,\eta)_x} [d\mu_x^\lambda((\xi) \times d\mu_x^\lambda(\eta) \times dt].$$

Consider $\widetilde{w} = \widetilde{w}(\xi, \eta, t)$ close to $S_x\widetilde{M}$ and write the coordinates (v, v', t) of $w = p\,\widetilde{w}$ as:

$$v = p\,(\pi_x^{-1}(\eta)), \quad v' = p\,(W^{ss}(\pi_x^{-1}(\eta)) \cap \gamma_{[\xi,\eta]}), \quad t = t.$$

In particular, w is close to v and

$$\theta_x^\lambda(\xi, \eta) = \theta_\lambda(w) k_\lambda(x, p(w), \xi) k_\lambda(x, p(w), \eta) \sim^{(1+\varepsilon)^2} \theta_\lambda(w) \sim^{(1+\varepsilon)^3} \theta_\lambda(v),$$

and

$$e^{-P(\lambda)(\xi,\eta)_x} \sim^{(1+\varepsilon)^2} 1.$$

We see that the measure $\widetilde{m}_\lambda$ has a density $\sim^{(1+\varepsilon)^8} \Omega_\lambda \theta_\lambda^{-2}(v)$ with respect to the product measure $d\mu_x^\lambda(\xi) \times d\mu_x^\lambda(\eta) \times dt$. The mapping $\xi \mapsto v'$ sends the measure $d\mu_x^\lambda$ to a measure with density $\sim^{(1+\varepsilon)^4} 1$ with respect to the measure $d\mu_\lambda^{ss}(v')$. This implies that we can write, in the neighborhood of $S_{px}M$, the measure m_λ in the coordinates (v, v', t) as

$$dm_\lambda(v, v', t) \sim^{(1+\varepsilon)^{12}} \Omega_\lambda \theta_\lambda^2(v)[d\mu_\lambda^{uu}(v) \times d\mu_\lambda^{ss}(v') \times dt].$$

Since $\delta'(1+\varepsilon)^{-1} \le \int \chi_-\left(\frac{t}{\delta'}\right) dt \le \int \chi_+\left(\frac{t}{\delta'}\right) dt \le \delta'(1+\varepsilon)$, it follows that

$$\begin{aligned}
\int F_\lambda^+(w, \lambda) dm_\lambda &\le (1+\varepsilon)^{12} \Omega_\lambda \int \chi_\pm\left(\frac{t}{\delta'}\right) dt \int_{S_{px}M} \left(\int_{W^{ss}_{loc}(v)} a_\lambda^\pm(v') d\mu_\lambda^{ss}(v') \right) f(v) d\mu_\lambda^{uu}(v) \\
&\le (1+\varepsilon)^{14} \Omega_\lambda \delta' \beta_0 \int_{S_{px}M} f(v) d\mu_\lambda^{uu}(v),
\end{aligned}$$

and

$$\int F_\lambda^-(w, \lambda) dm_\lambda \ge (1+\varepsilon)^{-14} \Omega_\lambda \delta' \beta_0 \int_{S_{px}M} f(v) d\mu_\lambda^{uu}(v).$$

Similarly, in the δ'-neighborhood of any lift of $S_{py}M$, we have, in the (u, u', s) coordinates, $u \in S_y\widetilde{M}, u' \in W^{uu}_{loc}(u), |s| \le 2\delta'$,

$$dm_\lambda(u, u', s) \sim^{(1+\varepsilon)^{12}} \Omega_\lambda \theta_\lambda^2(u)[d\mu_\lambda^{uu}(u') \times d\mu_\lambda^{ss}(u) \times ds].$$

The analog computation yields that

$$\begin{aligned}(1+\varepsilon)^{-14}\Omega_\lambda\gamma\beta_0'\int_{S_{py}M} h(-u)d\mu_\lambda^{ss}(u) \ &\leq\ \int H_\lambda^- dm_\lambda \\ \leq\ \ \int H_\lambda^+ dm_\lambda\ &\leq\ (1+\varepsilon)^{14}\Omega_\lambda\gamma\beta_0'\int_{S_{py}M} h(-u)d\mu_\lambda^{ss}(u).\end{aligned}$$

This shows Properties (3) and (4). Property (2) follows as $\int f d\mu_\lambda^{uu}$ and $\int h d\mu_\lambda^{ss}$ are bounded away from 0, uniformly in x, y and $\lambda \in [0, \lambda_0]$ by Corollary 2.8.

Step 4. Property (5)

Suppose $F_\lambda^-(w)H_\lambda^-(g_R w) \neq 0$. Set $w = (v, v', t)$ and $g_R w = (u, u', s)$, where $v \in$ *supp* f, $u \in$ *supp* h, v' is in the $r_\lambda^\pm(v)$-neighborhood of v in $W^{ss}(v)$ and u' is in the $r_\lambda^\pm(u)$-neighborhood of u' in $W^{uu}(u)$ with $|t| < \delta'/2$ and $|s| < \gamma < \delta'\epsilon/2$.

Since w is close to x and $g_R w$ is close to y, the geodesic from x to y remains close to the geodesic segment $\{g_t w : 0 \leq t \leq R\}$. In other words, there is a unique $v_0 \in S_{px}M$ such that $\{g_t v_0 : 0 \leq t \leq r\}$ remains close to $\{g_t w : 0 \leq t \leq R\}$.



FIGURE 2. M near x and y

Thus

$$\int F_\lambda^\pm(w)H_\lambda^\pm(g_R w)dm_\lambda = \sum_{(v_0,r)}\int_{\{g_t w\}_0^R \text{near} \{g_t v_0\}_0^r} \theta_\lambda^{-2} f(v)\theta_\lambda^{-2} h(-u)\chi_\pm(t/\delta')\chi_\pm(s/\gamma)a_\pm(v')b_\pm(u')dm_\lambda$$

By triangular inequalities on points in Figure 2,

$$|R-r| < |s-t| + \delta'\varepsilon < (1+3\varepsilon/2)\delta'.$$

Now fix such (v_0, r) and calculate the contribution of $\int F_\lambda^\pm(w)H_\lambda^\pm(g_R w)$ for w close to v_0.

Consider the closed path in SM near y in Figure 3 consisting of vectors

$$g_R v_0, g_R v, g_R v', g_R w, u', u, g_r v_0.$$

Then

$$\begin{aligned} d(g_R v_0, g_R v) &\leq d(g_R v, g_R v') + |t| + |s| + d(u', u) + d(u, g_r v_0) + d(g_r v_0, g_R v_0) \\ &\leq e^{-Ra_0} r_\lambda^+(v) + |t| + |s| + r_\lambda^+(u) + diam(S_{py}M) + |R-r| \\ &\leq e^{-Ra_0}\varepsilon\delta'/2 + \delta' + \gamma + \varepsilon\delta'/2 + C + \delta'(1+\varepsilon) \leq C \end{aligned}$$



FIGURE 3. SM near y

It follows that the distance between v_0 and v along the unstable direction is bounded above by $e^{-a_0R}C$. Since $S_{px}M$ is ε_1-transversal to $W^{ss}(v)$ (for some ε_1 independent of v),

$$d(v_0, v) < e^{-a_0R}C'.$$

Let us choose R_0 large enough so that if $R \geq R_0$, then $d(v_0, v) \leq \delta'$.



FIGURE 4. SM near x

Similarly, considering a closed path in the Figure 4, $d(g_{r-R}v_0, g_{-R}u) < C$, thus

$$d(g_r v_0, u) < e^{-Rh}C' < \delta'.$$

Choose another system of coordinates $W^{uu} \times W^{ss} \times \mathbb{R}$ for w near v_0 as follows :

$$w = (u'', v'', t),$$

where u'' and v'' are the projections of v' on $W^{uu}(v_0)$ and $W^{ss}(v_0)$, respectively, and t is defined as above. Since v, v' are δ'-close to u'', v'', respectively, the measure m_λ satisfies

$$dm_\lambda(u'', v'', t) \sim^{(1+\varepsilon)^{16}} \Omega_\lambda \theta_\lambda^2(v)[d\mu_\lambda^{uu}(u'') \times d\mu_\lambda^{ss}(v'') \times dt]. \tag{4.4}$$

Note that, for fixed v, v', u, u', the distances $d(\pi v, \pi u) = r$ and $d(\pi v', \pi u') = s + R - t$ are fixed. We are reduced to compute

$$\Omega_\lambda \int\int \left[\int \chi_\pm(t/\delta')\chi_\pm(s/\gamma)dt\right] a_\pm(v')b_\pm(u')f(v)\theta_\lambda^{-2}h(-u)d\mu_\lambda^{ss}(v'')d\mu_\lambda^{uu}(u'').$$

Let us first estimate the integral with respect to t. From $F_\lambda^-(w)H_\lambda^-(g_R w) \neq 0$, by triangular inequality,

$$d(v', g_{-R}u') < d(v', w) + d(w, g_{-R}u') \leq \delta'/2(1+\varepsilon) + \delta'\varepsilon/2(1+\varepsilon) = \delta'/2.$$

Clearly $\int \chi_-(t/\delta')\chi_-(s/\gamma)\,dt \leq \int \chi_+(s/\gamma)ds = (1+\varepsilon)^2\gamma$.

On the other hand, $d(v', g_{-R}u') < \delta'/2$ implies that

$$\int \chi_+(t/\delta')\chi_+(s/\gamma)\,dt \geq \int \chi_-(s/\gamma)ds = (1+\varepsilon)^{-2}\gamma.$$

Since v'' is δ'-close to v', we have

$$\int a_\pm(v')\,d\mu_\lambda^{ss}(v'') \ \sim^{(1+\varepsilon)^4} \int a_\pm(v')\,d\mu_\lambda^{ss}(v').$$

Indeed, the change of variables from v'' to v' is done by sliding along the central unstable manifolds between two close pieces of strong stable manifolds and the Radon-Nikodym derivative is

$$k_\lambda^2(v'', v', \cdot)e^{-P(\lambda)b(v'',v',\cdot)} \ \sim^{(1+\varepsilon)^4} 1.$$

We obtain, independently of u' and t that

$$\int a_-(v')\,d\mu^{ss}(v'') \ \leq \ (1+\varepsilon)^5\beta_0, \quad (1+\varepsilon)^{-5}\beta_0 \ \leq \ \int a_+(v')\,d\mu^{ss}(v'').$$

In the same way we have, independently of v'', t:

$$\begin{aligned} \int_{W_{loc}^{uu}(v_0)} b_\pm(u')\,d\mu_\lambda^{uu}(u'') \quad &= \quad \int_{W_{loc}^{uu}(v_0)} b_\pm(u')d\mu_\lambda^{uu}(u')\,\frac{d\mu_\lambda^{uu}(u'')}{d\mu_\lambda^{uu}(u')} \\ &\sim^{(1+\varepsilon)^4} \quad \frac{d\mu_\lambda^{uu}}{dg_{-r}\mu_\lambda^{uu}}(v_0)\int_{W_{loc}^{uu}(v_0)} b_\pm(u')d\mu_\lambda^{uu}(u'), \end{aligned}$$

since u'', u' are δ'-close to v_0 and $g_r v_0$, respectively. Altogether, we get for each (v,t) with $v \in S_{px}M \cap g_{-t}S_{py}M$ for some $t, |t-R| \leq \delta'/2$ a contribution $C_\pm(v,t)$ to the integral of $F_\lambda^- H_- \circ g_R, F_+H_+ \circ g_R$ satisfying:

$$(1+\varepsilon)^{-32}C_-(v,t) \ \leq \Omega_\lambda\gamma'\beta_0\beta_0' f(v)(\theta_\lambda^{-2}h)(-g_R v)\frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) \ \leq (1+\varepsilon)^{32}C_+(v,t). \tag{4.5}$$

Summing all these contributions for all relevant pairs (v,t), we obtain Property (5).

Step 5. End of the proof of Proposition 4.4

By Properties (1), (2) we can apply Proposition 4.1 and find R_0, δ_0 independent of λ, x, y such that for $R > R_0, \lambda \in [\lambda_0 - \delta_0, \lambda_0]$,

$$\begin{aligned} \int F_\lambda^- H_- \circ g_R\,dm_\lambda \ &\sim^{(1+\varepsilon)} \ \int F_\lambda^-\,dm_\lambda \int H_-\,dm_\lambda \\ \int F_+H_+ \circ g_R\,dm_\lambda \ &\sim^{(1+\varepsilon)} \ \int F_+\,dm_\lambda \int H_+\,dm_\lambda. \end{aligned}$$

We get

$$\begin{aligned}
&\Omega_\lambda \gamma \beta_0 \beta_0' \sum_{\{(v,t): v \in S_{px}M \cap g_{-t} S_{py}M\}} \chi_{\delta'}(R-t) f(v) (\theta^{-2} h)(-g_t v) \frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v) \\
\sim^{(1+\varepsilon)^{61}} \quad &\Omega_\lambda^2 \delta' \beta_0 \gamma \beta_0' \int_{S_{px}M} f(v) d\mu_\lambda^{uu}(v) \int_{S_{py}M} h(-v) d\mu_\lambda^{ss}(u),
\end{aligned}$$

which the statement of Proposition 4.4 after dividing both members by $\Omega_\lambda \gamma \beta_0 \beta_0'$. □

4.2. **Convergence of measures.** We state in this subsection several consequences and variants of Theorem 4.3 which will be used in the next Sections. Set $\Omega := \Omega_{\lambda_0}$ and $\Upsilon := \Upsilon_{\lambda_0}$.

First, observe that the expression (4.1) is continuous in λ as $\lambda \to \lambda_0$ by Corollary 3.13. By choosing $\delta_1 = \delta_1(f, h, \varepsilon)$ such that for $\lambda \in [\lambda_0 - \delta_1, \lambda_0]$

$$\begin{aligned}
&\Omega \int_{\partial \widetilde{M}} f(p \circ \pi_x^{-1} \xi) d\mu_x^{\lambda_0}(\xi) \int_{M_0} \left(\int_{\partial \widetilde{M}} h(p \circ \pi_y^{-1} \xi) d\mu_y^{\lambda_0}(\xi) \right) \\
\sim^{(1+\varepsilon)} \quad &\Omega_\lambda \int_{\partial \widetilde{M}} f(p \circ \pi_x^{-1} \xi) d\mu_x^{\lambda}(\xi) \int_{M_0} \left(\int_{\partial \widetilde{M}} h(p \circ \pi_y^{-1} \xi) d\mu_y^{\lambda}(\xi) \right),
\end{aligned}$$

we obtain, with the notations of Theorem 4.3,

Corollary 4.6. *Given $\varepsilon' > 0$ and positive Hölder continuous functions f, h on SM, there is $R(f, h, \varepsilon')$ and $\delta(f, h, \varepsilon')$ such that if $R > R(f, h, \varepsilon')$ and $\lambda_0 - \lambda < \delta(f, h, \varepsilon')$, for all $x \in \widetilde{M}$,*

$$\begin{aligned}
&e^{-RP(\lambda)} \int_{S(x,R)} f(p\, v_x^y) h(p\, v_y^x) G_\lambda^2(x, y) dy \qquad \sim^{(1+\varepsilon')^4} \\
&\Omega \int_{\partial \widetilde{M}} f(Dp \circ \pi_x^{-1} \xi) d\mu_x^{\lambda_0}(\xi) \int_{M_0} \left(\int_{\partial \widetilde{M}} h(p \circ \pi_y^{-1} \xi) d\mu_y^{\lambda_0}(\xi) \right) dVol(y).
\end{aligned}$$

We have to take $\delta(f, h, \varepsilon') < \delta_1(f, h, \varepsilon')$ and apply Theorem 4.3.

Corollary 4.7. *Fix $x \in \widetilde{M}$. Given $\varepsilon' > 0$ and a positive Hölder continuous functions f on $S_x\widetilde{M}$, there is $R(f, \varepsilon')$ and $\delta(f, \varepsilon')$ such that if $R > R(f, \varepsilon')$ and $\lambda_0 - \lambda < \delta(f, \varepsilon')$,*

$$e^{-RP(\lambda)} \int_{S(x,R)} f(v_x^y) G_\lambda^2(x, y) dy \sim^{(1+\varepsilon')^4} \Omega \int_{\partial \widetilde{M}} f(\pi_x^{-1} \xi) d\mu_x^{\lambda_0}(\xi). \tag{4.6}$$

In particular, for $\lambda = \lambda_0$,

$$\lim_{R \to \infty} e^{-RP(\lambda_0)} \int_{S(x,R)} f(v_x^y) G_{\lambda_0}^2(x, y) dy = \Omega \int_{\partial \widetilde{M}} f(\pi_x^{-1} \xi) d\mu_x^{\lambda_0}(\xi).$$

Proof. Extend f to a Γ-invariant Hölder continuous function on $S\widetilde{M}$ and consider the function induced on SM. The statement follows by letting $h = 1$ in Corollary 4.6. □

Letting $f = 1$ in Corollary 4.6, we obtain the convergence announced in the introduction.

Corollary 4.8. *Fix $x \in \widetilde{M}$. As $R \to \infty$ and $\lambda \to \lambda_0$, the measures $m_{x,\lambda,R}$ of the introduction converge to the measure $\Omega \mu_x^{\lambda_0}(\partial \widetilde{M}) \overline{m}$ on SM, where $\overline{m}$ is given by, for any continuous function h on $C(SM)$,*

$$\int_{SM} h \, d\overline{m} \; = \; \int_{M_0} \left(\int_{\partial \widetilde{M}} h(p \circ \pi_y^{-1} \xi) d\mu_y^{\lambda_0}(\xi) \right) d\,Vol(y).$$

Observe also that in the proof of Theorem 4.3, the choice of $\delta(f, h, \varepsilon')$ is only made in Step 5, when we want to use the uniform mixing of Proposition 4.1. For a fixed λ, we can use instead the regular mixing of m_λ for Hölder continuous functions and obtain a proof of Proposition 2.10. We can write, taking $f = h = 1$:

Corollary 4.9. *In Proposition 2.10, the limit $D(x, \lambda)$ is given by*

$$D(x, \lambda) \; = \; \Omega_\lambda \, \mu_x^\lambda(\partial \widetilde{M}) \int_{M_0} \int_{\partial \widetilde{M}} d\mu_y^\lambda(\xi) d\,Vol(y) = \Omega_\lambda \, \mu_x^\lambda(\partial \widetilde{M}).$$

As a Corollary of the proof of Theorem 4.3 and Corollary 4.6, we state a generalization which will be needed in Section 6.1.

Proposition 4.10. *Given $\varepsilon > 0$ and positive Hölder continuous functions f, u on SM, there is $R(f, u, \varepsilon)$ and $\delta(f, u, \varepsilon)$ such that if $R > R(f, u, \varepsilon)$ and $\lambda_0 - \lambda < \delta(f, u, \varepsilon)$,*

$$\begin{aligned} & e^{-RP(\lambda)} \int_{S(x,R)} f(p\, v_x^z) \left(\frac{1}{R} \int_0^R u(g_s p\, v_x^z) ds \right) G_\lambda^2(x, y) dy \\ \sim^{1+\varepsilon} \quad & \Omega \int_{\partial \widetilde{M}} f(p \pi_x^{-1} \xi) d\mu_x^{\lambda_0}(\xi) \int_{SM} u dm_{\lambda_0}. \end{aligned}$$

By the same argument as in the proof of Theorem 4.3 and Corollary 4.6, we are reduced to show:

Proposition 4.11. *Given $\varepsilon' > 0$ and positive Hölder continuous functions f, u, there exist $R_1 = R_1(f, u, \varepsilon')$, $\delta_1(f, u, \varepsilon') > 0$ and $\delta', 0 < \delta' < \delta_0'$, such that for $R \geq R_1$, all $x, y \in M$ and all $\lambda \in [\lambda_0 - \delta_1, \lambda_0]$,*

$$\begin{aligned} & \sum_{\{(v,t) : v \in S_{px} M \cap g_{-t} S_{py} M\}} \chi_{\delta'}(R - t) f(v) \left(\frac{1}{R} \int_0^R u(g_s v) ds \right) (\theta^{-2})(-g_t v) \frac{d\mu_\lambda^{ss}}{dg_t \mu_\lambda^{ss}}(g_R v) \\ & \sim^{1+\varepsilon'} \; \delta' \Omega_\lambda \left(\int_{\partial \widetilde{M}} f(p \pi_x^{-1} \xi) \, d\mu_x^\lambda(\xi) \right) \left(\int_{\partial \widetilde{M}} d\mu_y^\lambda(\xi) \right) \int_{SM} u dm_\lambda. \end{aligned}$$

Proof. We choose the same ε such that $(1+\varepsilon)^{62} \leq 1+\varepsilon'$. We choose $\delta_1' < \delta'$ small enough that, for all $t > 0$, if $v, w \in SM$ are such that $d(v, w) < \delta_1'$ and $d(g_t v, g_t w) < \delta_1'$, then

$$\int_0^t u(g_s v) ds \; \sim^{1+\varepsilon} \; \int_0^t u(g_s w) ds.$$

This is possible because u is Hölder continuous and the two geodesics $g_s v, g_s w$ satisfy

$$d_{SM}(g_s v, g_s w) \ \leq \ C\delta_1' \max\{e^{-a_0 s}, e^{a_0(s-t)}\},$$

where C, a_0 are positive geometric constants. We then construct $F_\lambda^\pm, H_\lambda^\pm$ in the same way, with this new δ_1' (and accordingly possibly new $\beta_0, \beta_0', \gamma$). Properties (1) to (4) still hold. In the relation (4.5), we multiply the middle term by $\frac{1}{R}\int_0^R u(g_s v)ds$. We then loose one more factor $1+\varepsilon$ in relation (4.5) if we replace $C_\pm(v,t)$ by the corresponding contributions to the integrals

$$\int F_\lambda^\pm(w)\left(\frac{1}{R}\int_0^R u(g_s w)ds\right) H_\lambda^\pm(w)\, dm_\lambda(w).$$

For R large enough, the new Property (5) reads as:

$$\begin{aligned}
&\frac{(1+\varepsilon)^{-33}}{\Omega_\lambda \gamma\beta_0\beta_0'}\int F_\lambda^-(w)\left(\frac{1}{R}\int_0^R u(g_s w)ds\right) H_\lambda^-(g_R w) dm_\lambda(w)\\
\leq \quad & \sum_{\{(v,t): v\in S_{px}M\cap g_{-t}S_{py}M\}} \chi_{\delta'}(R-t) f(v)\left(\frac{1}{R}\int_0^R u(g_s v)ds\right)(\theta^{-2})(-g_t v)\frac{d\mu_\lambda^{uu}}{dg_{-t}\mu_\lambda^{uu}}(v)\\
\leq \quad & \frac{(1+\varepsilon)^{33}}{\Omega_\lambda \gamma\beta_0\beta_0'}\int F_\lambda^+(w)\left(\frac{1}{R}\int_0^R u(g_s w)ds\right) H_\lambda^+(g_R w) dm_\lambda(w).
\end{aligned}$$

We conclude as above, using Proposition 4.2 instead of Proposition 4.1. □

5. TOPOLOGICAL PRESSURE AT λ_0

In this section, we show that $P(\lambda_0) = 0$ and show direct consequences. We already know that $P(\lambda_0) \leq 0$ by Corollary 3.12. As in Section 7 in [**GL**], we show below in Proposition 5.1 that if $P(\lambda_0) < 0$, then $\int_{S(x,R)} G_\lambda^2(x,y)dy$ decays exponentially with R and conclude that $G_{\lambda_0+\varepsilon}(x,y)$ is finite, contradicting the definition of λ_0. Remark that instead of constructing a branching random walk as in [**GL**], we have a more direct argument.

Proposition 5.1. $P(\lambda_0) = 0$.

Proof. Assume that $P(\lambda_0) < 0$. We claim that for all $x \neq y$, there exists $\varepsilon > 0$ such that the function $\lambda \mapsto G_\lambda(x,y)$ admits a real analytic extension on an ε-neighborhood of λ_0. In particular, for $\lambda_0 < \lambda < \lambda_0 + \varepsilon$, the extension $G_\lambda(x,y)$ satisfies $G_\lambda(x,y) = \int_0^\infty e^{\lambda t}\wp(t,x,y)dt$, a contradiction with the definition of λ_0.

Let us now prove our claim. Fix $x \neq y \in \widetilde{M}$. By Proposition 2.2,

$$\frac{\partial^k}{\partial\lambda^k}G_\lambda(x,y) \ = \ k!\int_{\widetilde{M}^k} G_\lambda(x,x_1)G_\lambda(x_1,x_2)\cdots G_\lambda(x_k,y)\, d\mathrm{Vol}^k(x_1,x_2,\cdots,x_k).$$

The claim follows with $\varepsilon = 1/\rho$, if we show that there are positive numbers δ, C and ρ such that:

(5.1)
$$F_k \ := \ \int_{\widetilde{M}^k} G_{\lambda_0}(x,x_1)G_{\lambda_0}(x_1,x_2)\cdots G_{\lambda_0}(x_k,y)e^{\delta d(x,x_k)}\, d\mathrm{Vol}^k(x_1,x_2,\cdots,x_k) \ \leq \ C\rho^k.$$

Since $P(\lambda_0) < 0$, by Theorem 4.3, there is $C, \delta > 0$ such that, for all $x \in \widetilde{M}$, all $R > 1$,

$$\int_{S(x,R)} G^2_{\lambda_0}(x,z)\, dz \leq Ce^{-\delta R} \text{ and thus } \int_{\{y\in\widetilde{M}; d(x,y)\geq 2\}} G^2_{\lambda_0}(x,y)\, d\mathrm{Vol}(y) < +\infty.$$

By possibly choosing a smaller $\delta > 0$, we prove in the same way the following relation:

$$\int_{\{y\in\widetilde{M}; d(x,y)\geq 2\}} G^2_{\lambda_0}(x,y)e^{\delta d(x,y)}\, d\mathrm{Vol}(y) \ \leq \ B(x) \tag{5.2}$$

for some function $B(x)$. Being a Γ-invariant continuous function, $B(x)$ is bounded. For this choice of δ, we prove (5.1) by induction on k. For $k = 0$, (5.1) is trivial for a suitable choice of C. We are going to show that F_{k+1}/F_k is bounded independently of k (compare [**GL**] Proposition 4.7). We write:

$$F_{k+1} = \int_{\widetilde{M}}\int_{\widetilde{M}^k} G_{\lambda_0}(x,x_1)G_{\lambda_0}(x_1,x_2)\cdots G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y)e^{\delta d(x,z)}\, d\mathrm{Vol}^k(x_1,\cdots,x_k)d\mathrm{Vol}(z).$$

Relation (5.1) follows from Lemma 5.2.

Lemma 5.2. *There is $\rho > 0$ such that, for all $x_k \in \widetilde{M}$,*

$$\int_{\widetilde{M}} G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y)e^{\delta d(x,z)}\, d\,Vol(z) \ \leq \ \rho\, G_{\lambda_0}(x_k,y)e^{\delta d(x,x_k)}.$$

Proof. Let L be the geodesic segment going from y to x_k (the RHS is infinite if $x_k = y$). Let $pr(z)$ be the point of L realizing $d(z, pr(z)) = d(z, L)$.

If $d(z,L) \leq 2$, and $d(x_k,y) \geq 1, d(z,x_k) \geq 1, d(z,y) \geq 1,$ we have $G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y) \leq CG_{\lambda_0}(x_k,y)$ by Harnack inequality and the easy side of the Ancona inequality.
If $d(x_k,y) \geq 1$ and one of $d(z,x_k), d(z,y) \leq 1$, we also have this inequality by Harnack inequality and Proposition 2.1. On the other hand, it is clear that

$$\int_{\{z:d(z,L)\leq 2\}} e^{\delta d(x,z)} d\mathrm{Vol}(z) \ \leq \ C'e^{\delta d(y,x_k)} \ \leq C'e^{\delta d(x,y)}e^{\delta d(x,x_k)}.$$

If $d(z,L) \leq 2$, and $d(x_k,y) \leq 1$, we have

$$\int_{B(x_k,4)} G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y)\, d\mathrm{Vol}(z) \ \leq \ CG_{\lambda_0}(x_k,y)$$

by Corollary 3.18. Clearly, in this case, $d(x,z) \leq d(x,x_k) + 3$. It follows that

$$\int_{\{z:d(z,L)\leq 2\}} G_{\lambda_0}(x_k,z)G_{\lambda_0}(z,y)e^{\delta d(x,z)}\, d\mathrm{Vol}(z) \ \leq \ C\, G_{\lambda_0}(x_k,y)e^{\delta d(x,x_k)}.$$

For $d(z,L) \geq 2$, consider the thin geodesic triangles $(y,z,pr(z))$ and $(x_k,z,pr(z))$. The distance from $pr(z)$ to both geodesics $[z,y]$ and $[z,x_k]$ is bounded by the hyperbolicity constant. Using Harnack inequality and the hard side of the Ancona-Gouëzel inequality, we get

$$\begin{aligned} G_{\lambda_0}(x_k, z) &\leq CG_{\lambda_0}(x_k, pr(z))G_{\lambda_0}(pr(z), z) \\ G_{\lambda_0}(z, y) &\leq CG_{\lambda_0}(z, pr(z))G_{\lambda_0}(pr(z), y). \end{aligned}$$

Therefore, we have

$$\begin{aligned} & \int_{\{z\in\widetilde{M}:d(z,L)\geq 2\}} G_{\lambda_0}(x_k, z)G_{\lambda_0}(z, y)e^{\delta d(x,z)}\, d\mathrm{Vol}(z) \\ \leq\ & C\int_{\{z\in\widetilde{M}:d(z,L)\geq 2\}} G_{\lambda_0}(x_k, pr(z))G_{\lambda_0}(pr(z), y)e^{\delta d(x,pr(z))}G^2_{\lambda_0}(z, pr(z))e^{\delta d(z,pr(z))}\, d\mathrm{Vol}(z) \\ \leq\ & CG_{\lambda_0}(x_k, y)\int_{\{z\in\widetilde{M}:d(z,L)\geq 2\}} e^{\delta d(x,pr(z))}G^2_{\lambda_0}(z, pr(z))e^{\delta d(z,pr(z))}\, d\mathrm{Vol}(z). \end{aligned}$$

We use the function $\psi : \mathbb{R} \to \mathbb{R}, \psi(t) = \max(1-|t|, 0)$. Since $\int_{-1}^{d(y,x_k)+1} \psi(t-s)dt = 1$ for all s between 0 and $d(y, x_k)$, we obtain

$$\begin{aligned} & \int_{\{z\in\widetilde{M};d(z,L)\geq 2\}} e^{\delta d(x,pr(z))}G^2_{\lambda_0}(z, pr(z))e^{\delta d(z,pr(z))}\, d\mathrm{Vol}(z) \\ \leq\ & \int_{\{z\in\widetilde{M};d(z,L)\geq 2\}}\int_{-1}^{d(y,x_k)+1} \psi(t - d(y, pr(z)))e^{\delta d(x,pr(z)}G^2_{\lambda_0}(z, pr(z))e^{\delta d(z,pr(z))}\, dt\, d\mathrm{Vol}(z) \end{aligned}$$

Let γ_{y,x_k} be the geodesic through y and x_k (from y to x_k) with $\gamma_{y,x_k}(0) = y$. Set $w_t := \gamma_{y,x_k}(t)$ for t between -1 and $d(y, x_k) + 1$, by Fubini theorem, the right hand side of the previous inequality is equal to

$$\begin{aligned} & \int_{-1}^{d(y,x_k)+1}\int_{\{z\in\widetilde{M};d(z,L)\geq 2\}} \psi(t - d(y, pr(z)))e^{\delta d(x,pr(z))}G^2_{\lambda_0}(z, pr(z))e^{\delta d(z,pr(z))}\, d\mathrm{Vol}(z)\, dt \\ \leq\ & C\int_{-1}^{d(y,x_k)+1}\int_{\{z\in\widetilde{M};d(z,L)\geq 2\}} \psi(t - d(y, pr(z)))e^{\delta d(x,w_t)}G^2_{\lambda_0}(z, w_t)e^{\delta d(z,w_t)}\, d\mathrm{Vol}(z)\, dt \\ \leq\ & C\int_{-1}^{d(y,x_k)+1} e^{\delta d(x,w_t)}B(w_t)\, dt \\ \leq\ & C\int_{-1}^{d(y,x_k)+1} e^{\delta t}\sup_x B(x)\, dt\ \leq\ Ce^{\delta d(y,x_k)} \leq Ce^{\delta d(x,x_k)}, \end{aligned}$$

where the first inequality uses Harnack inequality, the second inequality uses (5.2) and the third inequality uses the fact that $d(x, w_t) \leq d(x, y) + t$. □

□

Proposition 5.3. *For* $x \neq y \in \widetilde{M}$,

$$\lim_{\lambda\to\lambda_0} -P(\lambda)\frac{\partial}{\partial\lambda}G_\lambda(x, y)\ =\ \Omega c(x, y),$$

where $c(x,y)$ is given by

$$c(x,y)=\int k_{\lambda_0}(x,y,\xi)d\mu_x^{\lambda_0}(\xi). \tag{5.3}$$

Moreover, the convergence is dominated for y varying in $B(x,1)$.

Proof. We have:

$$\begin{aligned}-P(\lambda)\frac{\partial}{\partial\lambda}G_\lambda(x,y) &= -P(\lambda)\int_{\widetilde{M}}G_\lambda(x,z)G_\lambda(y,z)d\mathrm{Vol}(z)\\ &= -P(\lambda)\int_0^\infty e^{P(\lambda)R}\left(\int_{S(x,R)}e^{-P(\lambda)R}k_\lambda(x,y,z)G_\lambda^2(x,z)dz\right)dR.\end{aligned}$$

By Corollary 3.18, for $d(x,y)\le 1$, $\int_{B(x,3)}G_\lambda(x,z)G_\lambda(y,z)d\mathrm{Vol}(z)\le CG_{\lambda_0}(x,y)$. As $P(\lambda)$ goes to 0, this part converges to 0. The convergence is dominated by Lemma 2.1.

On the other hand, as $R\to\infty$, the function $k_\lambda(x,y,z)$ is close to $k_\lambda(x,y,(v_x^z)^+)$ uniformly in λ (Theorem 1.4), thus it can be considered as a Hölder continuous function on S_xM. Observe that the constant $C(\max\{d(x,y),1\})$ in Proposition 3.9 is uniform for $d(x,y)\le 1$, so that the Hölder norm of $k_\lambda(x,y,(v_x^z)^+)$ is uniformly bounded for $\lambda\in[0\lambda_0]$ and $d(x,y)\le 1$. By Corollary 4.7, for R large enough and λ close enough to λ_0, uniformly for $d(x,y)\le 1$,

$$\int_{S(x,R)}e^{-P(\lambda)R}k_\lambda(x,y,z)G_\lambda^2(x,z)dz\sim^{1+\varepsilon}\Omega\int k_{\lambda_0}(x,y,\xi)d\mu_x^{\lambda_0}(\xi)=\Omega\, c(x,y). \tag{5.4}$$

As $\lambda\to\lambda_0$, $P(\lambda)\to P(\lambda_0)=0$, it follows that

$$\lim_{\lambda\to\lambda_0}-P(\lambda)\frac{\partial}{\partial\lambda}G_\lambda(x,y)=\lim_{\lambda\to\lambda_0,R\to\infty}\int_{S(x,R)}e^{-P(\lambda)R}k_\lambda(x,y,z)G_\lambda^2(x,z)dz=\Omega\, c(x,y).$$

Indeed, Let R',λ' be such that (5.4) holds for $R\ge R',\lambda'\le\lambda<\lambda_0$ (R' and λ' depend only on $\max\{d(x,y),1\}$). We cut the integral

$$-P(\lambda)\int_0^\infty e^{P(\lambda)R}\left(\int_{S(x,R)}e^{-P(\lambda)R}G_\lambda(y,z)G_\lambda(x,z)\,dz\right)dR.$$

into $-P(\lambda)\int_0^3, -P(\lambda)\int_3^{R'}, -P(\lambda)\int_{R'}^\infty$. The first integral converges to 0, and the convergence is dominated for y varying in $B(x,1)$, the second integral goes to 0 and the third satisfies (5.4). □

In particular, since Ω and $c(x,y)$ are positive numbers, $\frac{\partial}{\partial\lambda}G_\lambda(x,y)$ goes to infinity as $\lambda\to\lambda_0$.

Remark 5.4. It follows from the proof above that

$$\lim_{\lambda\to\lambda_0}-P(\lambda)\int_{\widetilde{M}\setminus B(x,1)}G_\lambda(x,z)G_\lambda(x,z)d\mathrm{Vol}(z)=\Omega\int k_{\lambda_0}(x,x,\xi)d\mu_x^{\lambda_0}(\xi)=\Omega\mu_x^{\lambda_0}(\partial\widetilde{M}).$$

Theorem 1.5 follows from Proposition 2.6, Corollary 3.13 and Proposition 5.1 by letting $\lambda \to \lambda_0$.

Finally with Corollary 4.7 ($f = 1$ for the first limit and $f = k(x, y, z)$ for the second limit), we obtain

Proposition 5.5. *For $x, y \in \widetilde{M}$, as $R \to \infty$, we have, with the above notations*

$$\int_{S(x,R)} G^2_{\lambda_0}(x,z)\,dz \;\to\; \Omega\mu_x^{\lambda_0}(\partial\widetilde{M}), \qquad \int_{S(x,R)} G_{\lambda_0}(x,z)G_{\lambda_0}(y,z)\,dz \;\to\; \Omega c(x,y),$$

and

$$\int_{S(x,R)} f(v_x^z)G^2_{\lambda_0}(x,z)\,dz \;\to\; \Omega\int_{\partial\widetilde{M}} f(p\,\pi_x^{-1}(\xi))\mu_x^{\lambda_0}(\xi).$$

Observe that the last limit can serve as another definition of the $\mu_x^{\lambda_0}$.

Let us now prove another characterization of $\mu_x^{\lambda_0}$ which is Proposition 1.6.

Proof of Theorem 1.6. Suppose that we have a Γ-equivariant family $\{\nu_x\}$ of measures which minimizes $\mathcal{E}(\nu)$. Mohsen showed that (Lemma 5 in [**Mo**]) for each x, ξ, the function $\sqrt{k(x,\cdot,\xi)}$ is a $(-\lambda_0)$-eigenfunction, thus by Theorem 1.4, it satisfies

$$\sqrt{k(x,y,\xi)} = \int k_{\lambda_0}(x,y,\eta)d\nu_{x,\xi}(\eta),$$

for some family $\{\nu_{x,\xi}\}$ of probability measures on $\partial\widetilde{M}$. Recall that cocyles of Γ-equivariant families $\{\nu_x\}$ and $\{\mu_x^{\lambda_0}\}$ are normalized [**L2**], i.e. the pressure satisfies

$$P(-\frac{d}{dt}\ln k(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t), \widetilde{v}^+)\Big|_{t=0}) = 0,$$

and

$$P(-2\frac{d}{dt}\ln k_{\lambda_0}(\gamma_{\widetilde{v}}(0), \gamma_{\widetilde{v}}(t), \widetilde{v}^+)\Big|_{t=0}) = 0.$$

By definition, for any potential ϕ and $R > 0$, $P(\phi) = P(\frac{1}{R}\int_0^R \phi(g_t v))$. In particular, $P(K_R) = P(K_{R,0}) = 0$ for

$$K_R(x,\xi) = -\frac{1}{R}\ln k(x, \gamma_{\tilde{v}}(R), \xi) \text{ and } K_{R,0}(x,\xi) = -\frac{2}{R}\ln k_{\lambda_0}(x, \gamma_{\tilde{v}}(R), \xi),$$

where $\tilde{v} = \pi_x^{-1}(\xi)$. Suppose that there exist x, ξ such that $\nu_{x,\xi}(\eta) \neq \delta_{\xi,\eta}$. We claim that for R large enough, $K_R \geq K_{R,0}$ and $K_R(x,\xi) > K_{R,0}(x,\xi)$. Since the Gibbs measure m_{λ_0} is fully supported, it implies that $P(K_R) \geq h_{m_{\lambda_0}}(g) + \int_{SM} K_R dm_{\lambda_0} > P(K_{R,0}) = 0$, leading to a contradiction.

It remains to show that, for R large enough,

$$\int k_{\lambda_0}(x, \gamma_{\tilde{v}}(R), \eta)d\nu_{x,\xi}(\eta) \leq k_{\lambda_0}(x, \gamma_{\tilde{v}}(R), \xi)$$

and

$$\int k_{\lambda_0}(x, \gamma_{\tilde{v}}(R), \eta)d\nu_{x,\xi}(\eta) < k_{\lambda_0}(x, \gamma_{\tilde{v}}(R), \xi)$$

if $\nu_{x,\xi} < 1$. Let $\nu_{x,\xi} = a\delta_{\xi,\eta} + (1-a)\nu'_{x,\xi}$, where $\nu'_{x,\xi}(\{\xi\}) = 0$. Choose $r > 0$ such that $\nu'_{x,\xi}(B(\xi,r)) < 1/3C$, where C is the constant from Lemma 3.15. By Lemma 3.15, for x and $y = \gamma_{\tilde{v}}(R)$,

$$\int_{B(\xi,r)} k_{\lambda_0}(x,y,\eta)d\nu'_{x,\xi}(\eta) < 1/3k_{\lambda_0}(x,y,\xi).$$

On the complement of $B(\xi,r)$, we have

$$k_{\lambda_0}(x,y,\eta) \le k_{\lambda_0}(x,y,\xi)CG_{\lambda_0}(y,x_0)^2 \le k_{\lambda_0}(x,y,\xi)CC_r^2G_{\lambda_0}(x,y)^2.$$

Note that in the last inequality, we use Remark 3.16 to replace $G_{\lambda_0}(y,x_0)$ by $C_rG_{\lambda_0}(y,x)$ on $B(\xi,r)^c$. Thus we have

$$\int_{B(\xi,r)^c} k_{\lambda_0}(x,y,\eta)d\nu'_{x,\xi}(\eta) < k_{\lambda_0}(x,y,\xi)CC_r^2G_{\lambda_0}(x,y)^2.$$

If $a \neq 1$, then by Theorem 1.3, choose R large enough so that $G_{\lambda_0}(x,y)^2 < \frac{1}{3CC_r^2}$. It follows that

$$\int k_{\lambda_0}(x,y,\eta)d\nu_{x,\xi}(\eta) \le k_{\lambda_0}(x,y,\xi)\left(a + (1-a)\left(1/3 + CC_r^2G_{\lambda_0}(y,x)^2\right)\right) < k_{\lambda_0}(x,y,\xi).$$

We conclude that $\{\nu_x\}$ coincide with $\{\mu_x^{\lambda_0}\}$. □

6. Proof of Theorem 1.1

6.1. **Derivative of the Green function.** In this subsection, we establish

Theorem 6.1. *With the above notations, for $x \neq y \in \widetilde{M}$, as $\lambda \to \lambda_0$,*

$$\frac{\partial}{\partial\lambda}G_\lambda(x,y) \sim \frac{\sqrt{\Upsilon}}{2\sqrt{\lambda_0-\lambda}}c(x,y).$$

where $c(x,y)$ is given by (5.3) and $\Upsilon = \Upsilon_{\lambda_0}$, given by (2.11).

Theorem 6.1 follows from the following Proposition.

Proposition 6.2. *(i) For $x \neq y \in \widetilde{M}$, as $\lambda \to \lambda_0$,*

$$-P^3(\lambda)\frac{\partial^2}{\partial\lambda^2}G_\lambda(x,y) \text{ converges to } 2\Omega^3/\Upsilon c(x,y). \tag{6.1}$$

(ii) For all $x \neq y \in \widetilde{M}$,

$$\lim_{\lambda\to\lambda_0} -P^3(\lambda)\int_{\widetilde{M}}\int_{\widetilde{M}} G_\lambda(x,z)G_\lambda(z,w)G_\lambda(w,y)d\,Vol(w)d\,Vol(z) = \Omega^3/\Upsilon c(x,y). \tag{6.2}$$

Moreover, the convergence in (6.1) and (6.2) is dominated for y varying in $B(x,1)$.

Proof. (Compare with [**G1**] Section 3.6.) By Proposition 2.2, for $x \neq y$, (6.1) follows from (6.2). Firstly observe that, by Corollary 3.18 and Proposition 2.2,

$$\int_{B(x,1)}\int_{\widetilde{M}} G_\lambda(x,z)G_\lambda(z,w)G_\lambda(w,y)d\mathrm{Vol}(w)d\mathrm{Vol}(z) \le C\frac{\partial}{\partial\lambda}G_\lambda(x,y).$$

By Proposition 5.3, when multiplied by $P^3(\lambda)$, this term goes to 0 and the convergence is dominated on $B(x,1)$. We have

$$\begin{aligned} & -P^3(\lambda)\int_{\widetilde{M}\setminus B(x,1)}\int_{\widetilde{M}} G_\lambda(x,z)G_\lambda(z,w)G_\lambda(w,y)d\mathrm{Vol}(w)d\mathrm{Vol}(z) \\ = & \ -P^3(\lambda)\int_{\widetilde{M}\setminus B(x,1)} G_\lambda^2(x,z)\frac{G_\lambda(y,z)}{G_\lambda(x,z)}\left(\int_{\widetilde{M}}\frac{G_\lambda(z,w)G_\lambda(w,y)}{G_\lambda(y,z)}d\mathrm{Vol}(w)\right)d\mathrm{Vol}(z) \\ = & \ P^2(\lambda)\int_1^\infty Re^{P(\lambda)R}\left(\int_{S(x,R)} e^{-P(\lambda)R}G_\lambda^2(x,z)k_\lambda(x,y,z)\Psi_\lambda(x,y,z)dz\right)dR, \end{aligned}$$

where

$$\Psi_\lambda(x,y,z) = \frac{1}{d(x,z)}\left(-P(\lambda)\int_{\widetilde{M}}\frac{G_\lambda(z,w)G_\lambda(w,y)}{G_\lambda(y,z)}d\mathrm{Vol}(w)\right).$$

As in the proof of Proposition 5.3, as $\lambda\to\lambda_0$, $P(\lambda)\to 0$ and the above integral converges towards

$$\lim_{R\to\infty,\lambda\to\lambda_0}\int_{S(x,R)} e^{-P(\lambda)R}G_\lambda^2(x,z)k_\lambda(x,y,z)\Psi_\lambda(x,y,z)dz \tag{6.3}$$

if the limit exists uniformly. We study this limit. First we study $\Psi_\lambda(x,y,z)$.

Lemma 6.3. *There is a Hölder continuous positive functions u on SM such that for large $d(x,z)$ and λ close to λ_0,*

$$\Psi_\lambda(x,y,z) \ \sim^{1+\varepsilon} \ \Omega\frac{1}{d(x,z)}\int_0^{d(x,z)} u(g_s v_x^z)ds.$$

The estimate is uniform in $x, y, d(x,y)\le 1$.

Proof. For $w\in\widetilde{M}$, write $pr(w)$ for the projection of w on the geodesic segment from x to z and $s(w)$ for the distance $d(pr(w),x)$. Set $\psi(v,w) := \psi(d(pr(w),\pi(v))) = \max\{|1-d(pr(w),\pi(v))|, 0\}$. For a positive real number T, we denote

$$N_T(x) \ := \ \{w; w\in\widetilde{M}, s(w)\le T,\}, \quad N_T(z) \ := \ \{w; w\in\widetilde{M}, s(w)\ge d(x,z)-T.\}$$

We choose $K = K(x,y,\varepsilon)$ such that the angle between the vectors $v_{pr(w)}^x$ and $v_{pr(w)}^y$ is small enough if $s(w)\ge K$. We estimate

$$-P(\lambda)\int_{\widetilde{M}}\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)}d\mathrm{Vol}(w)$$

by decomposing $\widetilde{M}$ into $N_{K+1}(x)\setminus B(x,K')$, $N_{K+1}(x)\cap B(x,K')$, $N_2(z)\setminus B(z,K')$, $N_2(z)\cap B(z,K')$ and $\{w; w\in\widetilde{M}, K+1\le s(w)\le d(x,z)-2\}$ for a suitable K'.

The constant K' is chosen so that, if $w\in N_{K+1}(x)\setminus B(x,K')$, then there exists a point w' on the geodesic segment $[z,w]$ such that $d(w',y)\le C, d(w',w)\ge R_0$. Similarly, if $w\in N_2(z)\cap B(z,K')$, then there exists a point w' on the geodesic segment $[y,w]$ such that $d(w',z)\le C, d(w',w)\ge R_0$. the choice of K' is uniform on $d(x,y)\le 1$.

We use the Ancona inequality (3.4) to write for instance

$$\begin{aligned}
& -P(\lambda)\int_{N_2(z)\setminus B(z,K')}\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)}dw\\
= & -P(\lambda)\int_{N_2(z)\setminus B(z,K')}G_\lambda^2(w,z)\frac{G_\lambda(w,y)}{G_\lambda(z,y)G_\lambda(w,z)}dw\\
\leq & -P(\lambda)C\int_{N_2(z)\setminus B(z,K')}G_\lambda^2(w,z)\frac{G_\lambda(w,y)}{G_\lambda(w',y)G_\lambda(w,w')}dw\\
\leq & -P(\lambda)C\int_{\widetilde{M}\setminus B(z,K')}G_\lambda^2(w,z)dw
\end{aligned}$$

which is bounded by Remark 5.4. The argument is similar for $N_{K+1}(x)\setminus B(x,K')$.

For $d(w,x)\leq K'$, $\dfrac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)}\leq C(d(x,y))G_\lambda(x,w)$ and the integral is finite by (2.1). The argument is similar for the integral over $B(z,K')$.

It remains to estimate

$$\begin{aligned}
\int_K^{d(x,z)-1}\overline{u}_\lambda(g_sv_x^z)ds &\leq -P(\lambda)\int_{K+1}^{d(x,z)-2}\left(\int_{\widetilde{M}}\psi(s-s(w))\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)}d\mathrm{Vol}(w)\right)ds\\
&\leq \int_{K+1}^{d(x,z)-2}\overline{u}_\lambda(g_sv_x^z)ds
\end{aligned}$$

where $\overline{u}_\lambda$ is the function on $S\widetilde{M}$ given by

$$\begin{aligned}
\overline{u}_\lambda(v) &= -P(\lambda)\int_{\widetilde{M}}\psi(v,w)\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)}d\mathrm{Vol}(w)\\
&= -P(\lambda)\int_0^\infty e^{P(\lambda)R}u_{\lambda,R}(v)dR,
\end{aligned}$$

where

$$u_{\lambda,R}(v):=e^{-P(\lambda)R}\int_{S(\pi(v),R)}G_\lambda^2(\pi(v),w)\psi(v,w)\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)G_\lambda^2(\pi(v),w)}dw,$$

We are reduced to show that there is a Γ-invariant function $\overline{u}$ on $S\widetilde{M}$ such that $\overline{u}_\lambda(v)\to\Omega u(v)$ as $\lambda\to\lambda_0$.

By Proposition 3.10, given $\varepsilon>0$, our choice of $K>0$ is such that if $w\notin N_K(x)\cup N_1(z)$ and $pr(w)$ is far from w, then

$$\frac{G_\lambda(w,z)G_\lambda(w,y)}{G_\lambda(z,y)G_\lambda^2(\pi(v),w)}=\frac{\theta^\lambda_{\pi(v)}(w,z)\theta^\lambda_{\pi(v)}(w,y)}{\theta^\lambda_{\pi(v)}(y,z)}\sim^{1+\varepsilon}\frac{\theta^\lambda_{\pi(v)}(\zeta,v^+)\theta^\lambda_{\pi(v)}(\zeta,v^-)}{\theta^\lambda_{\pi(v)}(v^-,v^+)},$$

where $\zeta=(v^w_{\pi(v)})^+$ (see the Figure 5).

Extend the projection pr to the boundary $\partial\widetilde{M}$. Then for w such that $w\notin N_K(x)\cup N_1(z)$, if R is large enough, $\psi(v,w)\sim^{1+\varepsilon}\psi(v,\zeta)$. Also, the functions $d_{\pi(v)}(\zeta,v^\pm)$ are bounded away from 0 and the function $\theta^\lambda_{\pi(v)}(\zeta,v^+)\theta^\lambda_{\pi(v)}(\zeta,v^-)$ is uniformly Hölder and



Figure 5. Approximating by Naim kernels

bounded away from 0. The denominator $\theta_{\lambda_0}(v)$ is also Hölder and the approximation is uniformly Hölder continuous. By Corollary 4.7 centered at $\pi(v)$, there is $R(\varepsilon)$ and $\delta(\varepsilon)$ such that for $R \geq R(\varepsilon), \lambda \in [\lambda_0 - \delta(\varepsilon), \lambda_0]$, we have $u_{\lambda,R}(v) \ \sim^{1+\varepsilon} \Omega u(v)$, where

$$u(v) \ = \ \int_{\partial \widetilde{M}} \psi(v,\zeta) \frac{\theta_{\pi(v)}(\zeta, v^+)\theta_{\pi(v)}(\zeta, v^-)}{\theta(v)} d\mu_{\pi(v)}^{\lambda_0}(\zeta). \tag{6.4}$$

The function u given by (6.4) is clearly Γ-invariant. We still denote u the corresponding function on SM.

The Lemma follows, since $P(\lambda) \to 0$ and $\overline{u}_\lambda(v) = -P(\lambda) \int_0^\infty e^{P(\lambda)R} u_{\lambda,R}(v)\, dR$. □

Note that $\int_{SM_0} u(v) dm_\lambda(v)$ can be computed. It is given by

$$\begin{aligned}
& \int \int_{\partial \widetilde{M}} \psi(v,\zeta) \frac{\theta_{\pi(v)}(\zeta, v^+)\theta_{\pi(v)}(\zeta, v^-)}{\theta(v)} d\mu_{\pi(v)}^{\lambda}(\zeta) dm_\lambda(v) \\
= & \int \int_{\partial \widetilde{M}} \psi(v,\zeta) \frac{\theta_{\pi(v)}(\zeta, v^+)\theta_{\pi(v)}(\zeta, v^-)}{\theta(v)} d\mu_{\pi(v)}^{\lambda}(\zeta) \Omega_\lambda \theta^2(v) d\mu_{\pi(v)}^{\lambda}(v^-) d\mu_{\pi(v)}^{\lambda}(v^+) dt \\
= & \ \Omega_\lambda / \Upsilon_\lambda \int_{\partial \widetilde{M}} \int_{(v^-, v^+, t) \in SM_0} \psi(v,\zeta) dt d\rho_{\pi(v)}^{\lambda}(v^+, v^-, \zeta) \\
= & \ \Omega_\lambda / \Upsilon_\lambda \ \tau_{\pi(v)}^{\lambda}(S^2 M) = \Omega_\lambda / \Upsilon_\lambda.
\end{aligned}$$

We are reduced to understanding the limit

$$\lim_{R \to \infty, \lambda \to \lambda_0} \int_{S(x,R)} e^{-P(\lambda)R} G_\lambda^2(x,z) k_\lambda(x,y,z) \left(\frac{1}{R} \int_0^R u(g_s v_x^z) ds \right) dz.$$

As in the proof of Proposition 5.3, we can replace $k_\lambda(x,y,z)$ by $k_{\lambda_0}(x,y,\pi_x(v_x^z))$ for R sufficiently large and λ close to λ_0. By Proposition 4.10, for R large and $\lambda_0 - \lambda$ small,

$$\int_{S(x,R)} e^{-P(\lambda)R} G_\lambda^2(x,z) k_{\lambda_0}(x,y,\pi_x(v_x^z)) \left(\frac{1}{R} \int_0^R u(g_s v_x^z) ds \right) dz \sim \Omega^2 c(x,y) \int_{SM} u\, dm_{\lambda_0}$$

by (5.3). This achieves the proof of Proposition 6.2. ☐

Proof. (of Theorem 6.1). Set $F(\lambda) = \frac{\partial}{\partial\lambda} G_\lambda(x,y)$. By Proposition 5.3 and Proposition 6.2 (i),

$$\lim_{\lambda\to\lambda_0} -P(\lambda)F(\lambda) = \Omega\, c(x,y) \quad \lim_{\lambda\to\lambda_0} -P^3(\lambda)F'(\lambda) = 2\frac{\Omega^3}{\Upsilon}c(x,y).$$

It follows that $\dfrac{2F'(\lambda)}{F(\lambda)^3}$ converge towards $\dfrac{4}{\Upsilon}(c(x,y))^{-2}$. Since $F(\lambda)$ goes to ∞ as $\lambda \to \lambda_0$, we conclude that $F(\lambda) \sim \frac{\sqrt{\Upsilon}}{2}\frac{c(x,y)}{\sqrt{\lambda_0-\lambda}}$. This is Theorem 6.1. ☐

By Proposition 5.3 and Theorem 6.1, we obtain

Corollary 6.4. *As* $\lambda \to \lambda_0$,

$$-\frac{P(\lambda)}{\sqrt{\lambda_0-\lambda}} \to \frac{2\Omega}{\sqrt{\Upsilon}}.$$

Applying Proposition 6.2(ii) and Corollary 6.4, we get

Corollary 6.5. *For all* $x \neq y \in \widetilde{M}$,

$$\lim_{\lambda\to\lambda_0} (\lambda_0-\lambda)^{3/2} \int_{\widetilde{M}\times\widetilde{M}} G_\lambda(x,z)G_\lambda(z,w)G_\lambda(w,y)d\mathrm{Vol}(z)d\mathrm{Vol}(w) \;=\; \frac{\sqrt{\Upsilon}}{8}c(x,y).$$

Moreover, the convergence is dominated for y varying in $B(x,1)$.

6.2. **Proof of Theorem 1.1 and Theorem 1.7.** Let F be a nonnegative C^∞ function on $\widetilde{M}$, with compact support. Set μ_F for the spectral measure of F, i.e. the Borel finite measure on the spectrum $[0,+\infty)$ of $-\Delta-\lambda_0$ such that, for all $m \geq 0$,

$$\int_{\widetilde{M}} F(x)\Delta^m F(x)\, d\mathrm{Vol}(x) \;=\; \int_0^{+\infty} (-\varpi-\lambda_0)^m\, d\mu_F(\varpi).$$

The function

$$c_F(t) \;:=\; \int_{\widetilde{M}\times\widetilde{M}} e^{\lambda_0 t}\wp(t,x,y)F(x)F(y)\, d\mathrm{Vol}(x)d\mathrm{Vol}(y) = \int_0^{+\infty} e^{-\varpi t}\, d\mu_F(\varpi)$$

is nonincreasing in t. Write

$$\int_0^{+\infty} e^{-st}t^2 c_F(t)\, dt \;=\; \int_{\widetilde{M}\times\widetilde{M}}\int_0^\infty t^2 e^{(\lambda_0-s)t}\wp(t,x,y)dtF(x)F(y)\, d\mathrm{Vol}(x)d\mathrm{Vol}(y).$$

By the same direct computation as in the proof of Proposition 2.2, this is

$$2\int_{\widetilde{M}\times\widetilde{M}} G_{\lambda_0-s}(x,z)G_{\lambda_0-s}(z,w)G_{\lambda_0-s}(w,y)F(x)F(y)d\mathrm{Vol}(z)d\mathrm{Vol}(w)d\mathrm{Vol}(x)d\mathrm{Vol}(y).$$

By Corollary 6.5, we have, as $s \to 0$,

$$s^{3/2}\int_0^{+\infty} e^{-st}t^2 c_F(t)\, dt \to \frac{\sqrt{\Upsilon}}{4}\int_{\widetilde{M}\times\widetilde{M}} c(x,y)F(x)F(y)\, d\mathrm{Vol}(x)d\mathrm{Vol}(y).$$

By Hardy-Littlewood Tauberian Theorem ([**F**] p. 445), we have

$$\int_0^T t^2 c_F(t) \;\sim\; \frac{\sqrt{\Upsilon}}{4\Gamma(5/2)} T^{3/2} \int_{\widetilde{M}\times\widetilde{M}} c(x,y)F(x)F(y)\, d\mathrm{Vol}(x) d\mathrm{Vol}(y). \tag{6.5}$$

Now we claim that

$$c_F(t) \sim \frac{\sqrt{\Upsilon}}{2\sqrt{\pi} t^{3/2}} \int_{\widetilde{M}\times\widetilde{M}} c(x,y)F(x)F(y)\, d\mathrm{Vol}(x) d\mathrm{Vol}(y).$$

Indeed, by setting $CT^{3/2}$ to be the right hand side of the equation 6.5, since $C_F(t)$ is a non-increasing function of t,

$$3C\varepsilon T^{3/2}/2 \sim \int_T^{T(1+\varepsilon)} t^2 c_F(t)dt \le c_F(T) \int_T^{T(1+\varepsilon)} t^2 dt \sim c_F(T)\varepsilon T^3,$$

and

$$3C\varepsilon T^{3/2}/2 \sim \int_{T(1-\varepsilon)}^{T} t^2 c_F(t)dt \ge c_F(T) \int_{T(1-\varepsilon)}^{T} t^2 dt \sim c_F(T)\varepsilon T^3.$$

(See also [**GL**] Lemma 9.5.) Since $c(x,y) = \int k_{\lambda_0}(x,y)d\mu_x$, and $k_{\lambda_0}(x,y)$ is smooth as a $(-\lambda_0)$-eigenfunction of the Laplacian, the function $c(x,y)$ is smooth (see the paragraph after Proposition 3.10). We claim that $\wp(t,x,y)$ has bounded gradient. Indeed, if $u(x,t)$ is a solution of the heat equation on $\widetilde{M}$ with Ricci curvature bounded below, then by a sharp gradient estimate by Souplet and Zhang [**SZ**], on $\{(x,t) : y \in B(x_0, R/2), s \in [t_0 - T/2, t_0]\}$,

$$\frac{|\nabla u(x,t)|}{u(x,t)} \le C\left(\frac{1}{R} + \frac{1}{\sqrt{T}} + \sqrt{k}\right)\left(1 + \ln \frac{\max u(x,t)}{\min u(x,t)}\right),$$

where the maximum and minimum are taken on the set $\{(x,t) : x \in B(x_0,R), t \in [t_0 - T, t_0]\}$. Thus it remains to show that for some $T_0 >> 0$, $\frac{\max u(x,t)}{\min u(x,t)}$ is bounded above and below for $x \in B(x_0,1)$ and for all $t > T_0$. We use parabolic Harnack inequality: Note that

$$\wp(t,x,y) \ge \int_{B(x_0,1)} \wp(s-1,x,z)\wp(1,z,y)dz \ge C \int_{B(x_0,1)} \wp(s-1,x,z)dz,$$

where C is the minimum of $\wp(1,z,y)$.

Since $\wp(t,x,y)$ has bounded gradients, we can take functions F with compact support converging to $\{x_0\}$ that approximate the Dirac measure δ_{x_0} to get

$$\lim_{t\to\infty} t^{3/2} e^{\lambda_0 t} \wp(t,x_0,x_0) \;=\; \frac{\sqrt{\Upsilon}}{2\sqrt{\pi}} c(x_0,x_0).$$

We get the general case of $x_0 \neq x_1$ (and thus Theorem 1.1) in the same way by considering functions that approximate $\delta_{x_0} + \delta_{x_1}$.

Theorem 1.7 follows from the above equation and definition of $c(x,y)$ (5.3).

7. UNIFORM MIXING

In this section, we establish a uniform power mixing of the geodesic flow for Gibbs measures, when the potential varies in a neighbourhood of the space $\mathcal{K}_\alpha$ of functions which will be defined shortly. The proof combines the ideas from [**P1**] and [**P2**], with a slightly different framework. For the comfort of the reader, we recall the different steps in our notations.

7.1. **Uniform mixing and three-mixing.** Let $\mathbb{X} := (X, \mathcal{A}, m; \varphi_t, t \in \mathbb{R})$ be a system with one parameter group $\{\varphi_t, t \in \mathbb{R}\}$ of measurable transformations of the space $(X, \mathcal{A})$ preserving a probability measure m. For bounded measurable functions f, h, u we define the correlations functions for $s, t \geq 0$:

$$\begin{aligned}
\rho_{f,h,m}(t) &= \int f(x)h(\varphi_t x)dm(x) - \int f\,dm \int h\,dm \\
\rho_{f,u,h,m}(s,t) &= \int f(x)u(\varphi_s x)h(\varphi_{s+t}x)dm(x) - \int f\,dm \int u\,dm \int h\,dm \\
\overline{\rho}_{f,u,h,m}(t) &= \frac{1}{t}\int_0^t \left[\int f(x)u(\varphi_s x)h(\varphi_t x)dm(x)\right] ds - \int f\,dm \int u\,dm \int h\,dm
\end{aligned}$$

The system $\mathbb{X}$ is called *mixing* if $\lim_{t\to\infty} \rho_{f,h,m}(t) = 0$ for all bounded functions f, h, *3-mixing* if $\lim_{s,t\to\infty} \rho_{f,u,h,m}(s,t) = 0$ for all bounded functions f, u, h and *average 3-mixing* if $\lim_{t\to\infty} \overline{\rho}_{f,u,h,m}(t) = 0$ for all bounded functions f, u, h. It is a well-known open problem whether mixing implies 3-mixing. It is easy to see that mixing implies average 3-mixing.

Let us consider the rate of mixing. A system $\mathbb{X}$ is called power mixing for a class $\mathcal{K}$ of functions if for $f, h, k \in \mathcal{K}$, $\rho_{f,h,m}(t)$ decays polynomially (see Theorem 7.2 for a precise statement). Below, we will show a uniform version of power mixing of the geodesic flow for the class $\mathcal{K} = \mathcal{K}_\alpha$ which we define now.

Let $\alpha > 0$. We denote $\mathcal{K}_\alpha$ the space of functions f on X such that $\|f\|_\alpha < \infty$, where

$$\|f\|_\alpha := \sup_x |f(x)| + \sup_{x\neq y} \frac{|f(x) - f(y)|}{(d(x,y))^\alpha}.$$

Let $\varphi \in \mathcal{K}_\alpha$. There is a unique invariant probability measure m_φ such that:

$$h_{m_\varphi}(g_1) + \int \varphi\,dm_\varphi = P(\varphi) := \sup_{m;(g_t)_*m=m} \left\{h_m(g_1) + \int \varphi\,dm\right\},$$

where $h_m(g_1)$ denotes the measure theoretic entropy of the g-invariant probability measure m (see e.g. [**PP**]). The mapping $\varphi \mapsto m_\varphi$ is continuous from $\mathcal{K}_\alpha$ to the space of measures on X endowed with the weak* topology.

The following property is important in Dolgopyat's approach to the speed of mixing.

Definition 7.1. *A system $\mathbb{X}$ is topologically power mixing if there exists $t_0, \delta > 0$ such that for any r and $t > \max\{\frac{1}{r^\delta}, t_0\}$,*

$$g_t(B(x,r)) \cap B(y,r) \neq \emptyset.$$

We now establish a local uniform power mixing for topological power mixing Anosov flows, for Gibbs measures associated to potentials φ, and for functions in $\mathcal{K}_\alpha$. The mixing rate is uniform as we vary the potential φ in a small neighbourhood in $\mathcal{K}_\alpha$, for α sufficiently small.

Theorem 7.2. *Let $\mathbb{X}$ to be a topological power mixing Anosov flow. Let $\varphi_0 \in \mathcal{K}_\alpha$ be a potential. There exist $\varepsilon > 0$ and $C_0', c_0' > 0$ such that for all φ in the ε-neighborhood of φ_0 and all $f, u, h \in \mathcal{K}_\alpha$, we have, for all positive s, t:*

$$\left|\rho_{f,u,h,m_\varphi}(s,t)\right| \ \leq\ C_0' \|f\|_\alpha \|u\|_\alpha \|h\|_\alpha [(1+s)^{-c_0'} + (1+t)^{-c_0'}]. \tag{7.1}$$

Proposition 7.3. *Let $\mathbb{X}$ to be a topological power mixing Anosov flow. Let $\varphi_0 \in \mathcal{K}_\alpha$ be a potential. There exist $\varepsilon > 0$ and $C_0, c_0 > 0$ such that for all φ in the ε-neighborhood of φ_0 and all $f, h \in \mathcal{K}_\alpha$, we have, for all positive t:*

$$\left|\rho_{f,h,m_\varphi}(t)\right| \ \leq\ C_0 \|f\|_\alpha \|h\|_\alpha (1+t)^{-c_0}. \tag{7.2}$$

Although it appears as if Proposition 7.3 follows from Theorem 7.2, we will prove both results simultaneously.

Corollary 7.4. *Let $\mathbb{X}$ to be a topological power mixing Anosov flow and $\varphi_0 \in \mathcal{K}_\alpha$ be a potential. There exist $\varepsilon > 0$ and $C_0', c_0' > 0$ such that for all φ in the ε-neighborhood of φ_0 and all $f, u, h \in \mathcal{K}_\alpha$, we have, for all positive s, t:*

$$\left|\overline{\rho}_{f,u,h,m_\varphi}(t)\right| \ \leq\ C_0' \|f\|_\alpha \|u\|_\alpha \|h\|_\alpha (1+t)^{-c_0'}. \tag{7.3}$$

We assume now that the system $\mathbb{X}$ is the geodesic flow $g_t, t \in \mathbb{R}$ on the unit tangent bundle $X = SM$, where M is a closed negatively curved manifold.

Liverani proved exponential mixing for contact Anosov flows for the Liouville measure, which implies exponential mixing for the geodesic flow on manifolds of negative curvature for the Liouville measure [**Li**]. It implies that the geodesic flow is topologically power mixing. Thus we can apply the above theorems to the geodesic flow and the Gibbs measure associated to φ_{λ_0} to obtain Propositions 4.1 and 4.2.

7.2. **Proof of Theorem 7.2 and Proposition 7.3.** First, following Bowen and Ruelle [**B**], [**BR**], we can reduce the problem to the corresponding problem on suspended symbolic flows by introducing Poincaré sections for the flow with Markov property (see also [**PP**] Chapter 9 and Appendix III), in such a way that Hölder continuous functions on SM correspond to Hölder continuous functions on the symbolic system.

We may thus assume that there is a subshift of finite type (Σ, σ) and a positive α-Hölder continuous function τ on Σ such that the system $\mathbb{X}$ is the suspension flow $\sigma_t(x,r) = (x, r+t)$ on the set $\Sigma^\tau := \{(x,r) : x \in \Sigma, 0 \leq r \leq \tau(x)\}/[(x,\tau(x)) \sim (\sigma x, 0)]$. Let us denote by $[a_0, \cdots, a_k]$ the cylinder set $\{x : x_i = a_i, i = 0, \cdots, k\}$. Let us also define d_α on the space Σ_+ of one-sided sequences with the left-shift by $d_\alpha(x,y) = \alpha^k$, where k is the first index for which x_k, y_k are not equal. Let us denote by $\mathcal{K}_\alpha(\Sigma_+)$ the space of d_α-Lipschitz functions on the space Σ_+ of one-sided sequences. Let $\varphi \in \mathcal{K}_\alpha(\Sigma^\tau)$ be a potential function on Σ^τ.

We may assume that the function τ is a function on Σ_+ in the sense that $\tau(x) = \tau(y)$ if the points x and y in Σ have the same nonnegative coordinates. Moreover, by possibly modifying α, we may assume that the function τ is a d_α-Lipschitz function on Σ_+. We may also assume that the functions $\phi(x) := \int_0^{\tau(x)} \varphi(x, r)\, dr$ are d_α-Lipschitz functions on Σ_+ as well and that ϕ are normalized so that $\mathcal{L}_\phi 1 = 1$, where

$$\mathcal{L}_\phi F(x) := \sum_{y;\sigma y = x} e^{\phi(y)} F(y) \tag{7.4}$$

is the transfer operator associated to ϕ.

It follows from these assumptions that the map $\mathcal{T} : \varphi \mapsto \phi$ is continuous from $\mathcal{K}_\alpha(\Sigma^\tau)$ into $\mathcal{K}_\alpha(\Sigma_+)$. The equilibrium measure m_φ for the function φ is of the form

$$m_\varphi = \frac{1}{\int \tau\, d\nu_\phi} (\overline{\nu}_\phi \otimes dr)\big|_{\Sigma^\tau},$$

where $\overline{\nu}_\phi$ is the unique σ-invariant probability measure on Σ such that its projection ν_ϕ to Σ_+ satisfies, for all functions $F \in C(\Sigma_+)$,

$$\int \mathcal{L}_\phi F\, d\nu_\phi = \int F\, d\nu_\phi. \tag{7.5}$$

Let us denote $\phi_k(x) = \phi(x) + \phi(\sigma(x)) + \cdots + \phi(\sigma^k(x))$. We choose an ε_1-neighborhood of ϕ_0 so that there exists a constant $C_1 \geq 1$ with

$$\left| \frac{e^{\phi_k(x)}}{e^{\phi_k(y)}} - 1 \right| \leq C_1 \alpha^{-k} d_\alpha(x, y) \ \text{ and } \ C_1^{-1} \leq \frac{\nu_\phi[a_0, \cdots, a_{k-1}]}{e^{\phi_k(x)}} \leq C_1 \tag{7.6}$$

for all ϕ in ε_1-neighborhood of ϕ_0 and all $x, y \in [a_0, \cdots, a_{k-1}]$. With those choices, for all ϕ, 1 is an isolated eigenvalue of $\mathcal{L}_\phi$ with eigenfunction the constant 1 (see [**PP**] for this and all the above standard reductions). A ball of radius r in Σ^τ contains a cylinder of length $-C \ln r$ in Σ times an interval of length cr in the flow direction. Its image on the manifold contains a ball of radius r^D, for some D. Therefore, we may assume that the suspension flow $\mathbb{X}$ is topologicallly power mixing.

Remark 7.5. The constants C_1, C_6, C_7, γ_3 that appear in this section coincide with the constants C_1, C_6, C_7, γ_3 in [**Me**].

7.2.1. *Properties of the complex transfer operator.* In this subsection, we will denote the space of complex d_α-Lipschitz continuous functions on Σ_+ by $\mathcal{K}_\alpha(\Sigma_+)$ again. We define the complex transfer operator $\mathcal{L}_{\phi + s\tau}, s \in \mathbb{C}$ on $\mathcal{K}_\alpha(\Sigma_+)$ by

$$\mathcal{L}_{\phi+s\tau} F(x) := \sum_{y;\sigma y = x} e^{\phi(y) + s\tau(y)} F(y).$$

Following [**Me**], set $s = a + ib$.

Proposition 7.6. *(Lemma 3.5 of* [**Me**]*) Let $\mathbb{X}$ be a topological power mixing Anosov flow. Let φ_0 be a α-Hölder continuous function. There are $\varepsilon > 0, \theta > 0, C > 0$ such that* *for all* $s = a + ib$ *with* $|b| > 1$ *and* $|a| < C^{-1}|b|^{-\theta}$,

$$||(1 - \mathcal{L}_{\phi+s\tau})^{-1}||_\alpha \leq C|b|^\theta, \tag{7.7}$$

for all $\phi = \mathcal{T}(\varphi)$ *with* $||\varphi - \varphi_0||_\alpha < \varepsilon$.

Proof. As in [**Me**], we carry the calculations for $0 \leq a \leq 1$ and $b > 1$. They are analogous for $b < -1$ and $-1 \leq a \leq 0$. We find a neighborhood $\mathcal{U}$ of ϕ_0 where (7.7) holds and choose ε such that $||\varphi - \varphi_0||_\alpha < \varepsilon$ implies that $\phi = \mathcal{T}(\varphi) \in \mathcal{U}$. We first have the preliminary estimate of [**Me**] in a uniform way.

Lemma 7.7. *(Lemma 3.7 of* [**Me**]*) There exist* $C_6, C_7, \gamma_3, \varepsilon_2 > 0$ *such that for all* ϕ *in the* ε_2*-neighborhood of* ϕ_0,

1. $|\mathcal{L}_{\phi+ib\tau}|_\infty \leq 1$,
2. $||\mathcal{L}^n_{\phi+ib\tau}F||_\alpha \leq C_6\{b|F|_\infty + \alpha^n||F||_\alpha\}$ *for all* $n \leq 1$ *and* $F \in \mathcal{K}_\alpha(\Sigma_+)$,
3. $||\mathcal{L}^n_\phi F - \int_{\Sigma_+} F d\nu_\phi||_\alpha \leq C_7\gamma_3^n||F||_\alpha$ *for all* $n \leq 1$ *and* $F \in \mathcal{K}_\alpha(\Sigma_+)$.

Proof. Part (2) comes from the fundamental inequality, thus C_6 is uniform in ϕ. Part (3) comes from the spectral gap of $\mathcal{L}_\phi$ thus C_7 and γ_3 can be chosen uniformly in a neighbourhood of ϕ_0 (see e.g. Kato [**Ka**] Theorem IV.3.1). □

As in [**Me**], define

$$||f||_b := \max\left\{|f|_\infty, \frac{1}{2C_6 b}\sup_{x\neq y}\frac{|f(x) - f(y)|}{(d(x,y))^\alpha}\right\}.$$

Since one may assume that $2C_6b > 1$, we have

$$||F||_b \leq ||F||_\alpha \leq (2C_6b + 1)||F||_b,$$

which implies that $||\mathcal{L}||_\alpha/||\mathcal{L}||_b$ lies between $2C_6b + 1$ and $(2C_6b + 1)^{-1}$.

Let $M_bF = e^{-ib\tau}F \circ \sigma$.

Definition 7.8. *The operator* M_b *has no approximate eigenfunction if there exists* $N \in \mathbb{N}$ *such that for every triple* $(\theta \geq N, \beta > 0, C \geq 1)$, *there exists* $k = k(\theta, \beta, C)$ *such that for all* (b, ρ, F) *with* $|F| = 1$ *and* $|b| > k$,

$$|M_b^{\beta \ln |b|}F(y) - e^{i\rho}F(y)| \geq C|b|^{-\theta},$$

for some y.

Lemma 7.9. *[uniform version of Section 3.2 of* [**Me**]*] Consider the following conditions.*

1. M_b *has no approximate eigenfunction.*
2. *There exists* $\theta > 0, \varepsilon > 0$ *such that* $||(I - \mathcal{L}^{-1}_{\phi+ib\tau})||_b = O(|b|^\theta)$ *as* $|b| \to \infty$, *for all* $\phi \in B_\varepsilon(\phi_0)$.
3. *There exist* $\theta, \epsilon > 0, \varepsilon > 0$ *and* $C \geq 1$ *such that* $||(I - \mathcal{L}_{\phi+s\tau})^{-1}||_b \leq C|b|^\theta$ *for all* $s = a + ib$ *with* $|b| > 1$, $|a| < \epsilon|b|^{-\theta}$ *and all* $\phi \in B_\varepsilon(\phi_0)$.

With the above notations, (1) implies (2) and (2) implies (3).

Proof. See Section 3.2 of [**Me**]. Let ε_1 be a constant such that C_1 in equation (7.6) and α_1, α_2 in [**Me**] are uniform in ϕ in ε_1-neighborhood of ϕ_0. Now let $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$, where ε_2 is chosen as in Lemma 7.7. ☐

Topological power mixing of $\mathbb{X}$ implies that M_b has no approximate eigenfunction by Sections 3 and 5 of [**D2**], thus Proposition 7.6 follows. ☐

We also recall that, by mixing of the geodesic flow, $\|\mathcal{L}_{\phi+ib\tau}\|_\alpha < 1$ for $b \neq 0$ (see [**PP**] Proposition 6.2). It follows that:

Proposition 7.10. *There is $\delta = \delta_{\phi_0} > 0, \varepsilon > 0$ such that, for all ϕ in a ε-neighborhood of ϕ_0, the mapping $s \mapsto \sum_n \mathcal{L}^n_{\phi+s\tau}$ is meromorphic on V*

$$V := \{s = a + ib : |b| < 2,\ |a| < \delta\}$$

with a simple pole at $s = 0$. Moreover, for a function $K \in \mathcal{K}_\alpha(\Sigma_+)$, the residue at $s = 0$ of the meromorphic function $s \mapsto \sum_n \mathcal{L}^n_{\phi+s\tau} K$ (with values in $\mathcal{K}_\alpha$) is a constant function with value $\nu_\phi(K)$.

Proof. For a fixed ϕ , this follows from [**PP**], Proposition 6.2 and Theorem 10.2, with a fixed $\delta = \delta_\phi$. By [**Ka**] Theorem IV.3.1 and compactness of the closure $\overline{V}$, there is a neighborhood $\mathcal{U}_0$ of ϕ_0 such that for $\phi \in \mathcal{U}_0$, the rest of the spectrum of $\mathcal{L}_{\phi+s\tau}, s \in \overline{V}$, is separated from 1 with $\delta = \delta_{\phi_0}$. ☐

7.2.2. *One-sided smooth functions.* We start by proving Theorem 7.2 for a particular space of functions. For $\alpha > 0$ and $M \in \mathbb{N}$, let $\mathcal{K}^+_{\alpha,M}$ be the set of functions f on Σ^τ with the following properties:

- for all $x \in \Sigma$, $f(x, r) = 0$ for r outside the interval $[\frac{\inf\tau}{3}, \frac{2\inf\tau}{3}]$,
- for all $x \in \Sigma$, $r \mapsto f(x, r)$ is of class C^M,
- for all $r \in [\frac{\inf\tau}{3}, \frac{2\inf\tau}{3}]$, $x \mapsto f(x, r)$ depends only on the nonnegative coordinates of x and
- the functions $\frac{\partial^k f}{\partial r^k}(x, r)$, for $0 \le k \le M$ are α-Hölder continuous in $x \in \Sigma$ and continuous in r.

For $f \in \mathcal{K}^+_{\alpha,M}$, we denote $||f||_{\alpha,M} := \sup_{r,k\le M} ||\frac{\partial^k f}{\partial r^k}(., r)||_\alpha$. The heart of the proof uses the arguments of [**D2**] to establish:

Proposition 7.11. *Let $\varphi_0 \in \mathcal{K}_\alpha(\Sigma_+)$as above. There exist $\varepsilon, C, c > 0$ and M such that for all φ, $\|\varphi - \varphi_0\| < \varepsilon$, all $f, u, h \in \mathcal{K}^+_{\alpha,M}$, we have, for all positive t_1, t_2:*

$$\left|\rho_{f,u,h,m_\lambda}(t_1, t_2)\right| \le C\|f\|_{\alpha,M}\|u\|_{\alpha,M}\|h\|_{\alpha,M}[(1 + t_1)^{-c} + (1 + t_2)^{-c}]. \tag{7.8}$$

Proof. Choose ε so that Proposition 7.6 and Proposition 7.10 holds for all φ with $\|\varphi - \varphi_0\| < \varepsilon$. Fix f, u, h, φ and write $\rho(t_1, t_2)$ for $\rho_{f,h,g,m_\varphi}(t_1, t_2)$. Assume first that $\int f\, dm_\lambda = \int g\, dm_\lambda = 0$. We consider the Laplace transform

$$\widehat{\rho}(s_1, s_2) = \int_{\mathbb{R}_+\times\mathbb{R}_+} \rho(t_1, t_2)e^{-s_1t_1}e^{-s_2t_2}\, dt_1\, dt_2$$

which makes sense a priori for $a_j > 0$, where $s_j = a_j + ib_j, j = 1, 2$. The following computation is valid for $a_j > 0$ and will allow us to extend analytically $\widehat{\rho}(s_1, s_2)$ to a larger domain and deduce the decay of $\rho(t_1, t_2)$ as t_1, t_2 go to infinity.

Lemma 7.12. *Consider the Laplace transforms F, U and H of the functions f, u and h given by:*

$$F(x,s) = \int_{\mathbb{R}} e^{-sr} f(x,r)dr,\ U(x,s) = \int_{\mathbb{R}} e^{-sr} u(x,r)dr,\ G(x,s) = \int_{\mathbb{R}} e^{-sr} g(x,r)dv.$$

Then, we have, for $a_1, a_2 > 0$:

$$\widehat{\rho}(s_1, s_2) = \sum_{n,m} \int_{\Sigma} H(x, s_2) \mathcal{L}^m_{\phi+s_2\tau} \big[U(., s_1 - s_2) \mathcal{L}^n_{\phi+s_1\tau} F(.., -s_1)(.) \big] (x)\, d\nu_\phi(x).$$

Proof. We develop:

$$\begin{aligned} \widehat{\rho}(s_1, s_2) &= \int_{\mathbb{R}_+ \times \mathbb{R}_+} \int_{\Sigma^\tau} f(x,r) u(\sigma_{t_1}(x,r)) h(\sigma_{t_1+t_2}(x,r)) e^{-s_1 t_1} e^{-s_2 t_2}\, dm_\varphi(x,r)\, dt_1\, dt_2 \\ &= \sum_{n,m} \int_{\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+} \int_{\Sigma} f(x,r) u(\sigma^n x, r + t_1 - \tau^n(x)) h(\sigma^{n+m} x, r + t_2 + t_1 - \tau^{n+m}(x)) \\ &\qquad\qquad e^{-s_1 t_1} e^{-s_2 t_2}\, dr\, d\nu_\phi(x)\, dt_1\, dt_2, \qquad (*) \end{aligned}$$

where $\tau^n(x) := \sum_{k=0}^{n-1} \tau(\sigma^k(x))$. Observe that for all fixed positive n, m the integral in t_1, t_2, r is also an integral over $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$. Then using the variables $w = r + t_1 - \tau^n(x)$ and $z = w + t_2 - \tau^m(\sigma^n x)$, the integral (*) can be written as

$$(*) = \int_{\Sigma} H(\sigma^{n+m} x, -s_2) e^{s_2 \tau^m(\sigma^n x)} U(\sigma^n x, s_2 - s_1) e^{s_1 \tau^n(x)} F(x, s_1)\, d\nu_\phi(x).$$

Using now the invariance of ν_ϕ under $\mathcal{L}_\phi$ (7.5) and the fact that $\mathcal{L}^n(HK \circ \sigma^n)(x) = K(x)\mathcal{L}(H)(x)$, we obtain:

$$\begin{aligned} (*) &= \int_{\Sigma} H(\sigma^m x, -s_2) e^{s_2 \tau^m(x)} U(x, s_2 - s_1) \mathcal{L}^n_{\phi+s_1\tau} F(\cdot, s_1)(x)\, d\nu_\phi(x) \\ &= \int_{\Sigma} H(x, -s_2) \mathcal{L}^m_{\phi+s_2\tau} \big[U(\cdot, s_2 - s_1) \mathcal{L}^n_{\phi+s_1\tau} F(\cdot\cdot, s_1)(\cdot) \big] (x)\, d\nu_\phi(x). \end{aligned}$$

The Lemma follows for $a_j = \Re\, s_j > 0$. □

By Proposition 7.6 and our choice of ε, we conclude that there exist constants δ, β such that, for all $\varphi, \|\varphi - \varphi_0\|_\alpha < \varepsilon$, the series of operators $\sum_n \mathcal{L}^n_{\phi+s\tau}$ converges and is analytic on the region U

$$U \ :=\ \{s, s = a + ib; |b| > 1,\ |a| < \frac{2\delta}{|b|^{\beta/2}}\}$$

and that there is a constant D_0 such that, for $s \in U$,

$$\| \sum_n \mathcal{L}^n_{\phi+s\tau} \|_\alpha \ \leq\ D_0 |b|^{D_0}. \tag{7.9}$$

Moreover, by Proposition 7.10, there is $\delta > 0$ such that the series of operators $\sum_n \mathcal{L}^n_{\phi+s\tau}$ converges and is meromorphic on the region V, has a simple pole at 0 and has residue at 0 the projection on the constant function $\nu_\phi(.)$.

On the other hand, since f, u and h belong to $\mathcal{K}^+_{\alpha,M}$, the functions $s \mapsto F(\cdot, s), s \mapsto U(\cdot, s)$ and $s \mapsto H(., s)$ are analytic into $\mathcal{K}_\alpha(\Sigma_+)$. Moreover, the functions $\|F(\cdot, s)\|_\alpha$, $\|U(\cdot, s)\|_\alpha$ and $\|H(\cdot, s)\|_\alpha$ decay at infinity like $(|b|)^{-M}$ and

$$\nu_\phi(F(.,0)) \;=\; \int_{\Sigma_+} \left(\int_{\mathbb{R}} f(x,r)\,dr \right) d\nu_\phi(x) \;=\; \int_{\Sigma^\tau} f\,dm_\varphi \;=\; 0.$$

It follows that the function

$$J(x,s) := \sum_n \mathcal{L}^n_{\phi+s\tau} F(\cdot,s)(x)$$

is analytic from $U \cup V$ into $\mathcal{K}_\alpha$ and that its $\mathcal{K}_\alpha$-norm is bounded by $C\|f\|_{\alpha,M}(1+|b|)^{D_0-M}$ as $|b| \to \infty$. Summarizing, the function $\widehat{\rho}(s_1,s_2)$ admits an analytic extension to $\{(s_1,s_2); s_1 \in U \cup V, a_2 > 0\}$ and this extension satisfies:

$$\widehat{\rho}(s_1,s_2) \;=\; \sum_m \int_{\Sigma_+} H(x,-s_2)\mathcal{L}^m_{\phi+s_2\tau}[U(\cdot,s_2-s_1)J(\cdot,s_1)](x)\,d\nu_\phi(x).$$

As before, for each fixed $s_1 \in U \cup V$, the mapping $s_2 \mapsto \sum_m \mathcal{L}^m_{\phi+s_2\tau}[U(\cdot,s_2-s_1)J(\cdot,s)](x)$ is meromorphic from $U \cup V$ with a unique simple pole at $s_2 = 0$ and a residue a constant function on Σ_+ with value $C_0(s_1)$. Therefore, for all $s_1 \in U \cup V$, $s_2 \mapsto \widehat{\rho}(s_1,s_2)$ admits a meromorphic extension to $U \cup V$ of the form

$$\widehat{\rho}(s_1,s_2) \;=\; \frac{C_0(s_1)\int_{\Sigma_+} H(x,0)\,d\nu_\phi(x)}{2\pi i s_2} + \overline{\rho}(s_1,s_2),$$

where $\overline{\rho}(\xi,\eta)$ is an analytic function on $(U \cup V) \times (U \cup V)$ such that

$$|\overline{\rho}(s_1,s_2)| \;\le\; C\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}(1+|b_2|)^{D_0-M}(1+|b_1-b_2|)^{-M}(1+|b_1|)^{D_0-M}.$$

We again have $\int_{\Sigma_+} H(x,0)\,d\nu_\phi(x) = 0$ by our condition that $\int h\,d\mu_\phi = 0$ and finally, the function $\widehat{\rho}(s_1,s_2)$ admits an analytic extension to $(U \cup V) \times (U \cup V)$ and satisfies:

$$|\widehat{\rho}(s_1,s_2)| \;\le\; C\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}(1+|b_2|)^{D_0-M}(1+|b_1-b_2|)^{-M}(1+|b_1|)^{D_0-M}.$$

We now compute $\rho(t_1,t_2)$ as the Laplace inverse of $\widehat{\rho}(s_1,s_2)$ by integrating on the imaginary axis in s_2 and in s_1. For a fixed $s_1 \in U \cup V$, we can move the curve of integration in s_2 to the curve

$$\Gamma := \{-\delta \min\{1, \frac{1}{|b|^\beta}\} + ib; b \in \mathbb{R}\}.$$

We obtain that the function $\widetilde{\rho}(s_1,t_2)$

$$\begin{aligned} \widetilde{\rho}(s_1,t_2) &:= \frac{-1}{4\pi^2}\int_{\mathbb{R}} \widehat{\rho}(s_1,b_2)e^{ib_2t_2}\,db_2 \\ &= \frac{-1}{4\pi^2}\left(\int_{-1}^{+1} \widehat{\rho}(s_1,-\delta+ib_2)e^{ib_2t_2}e^{-\delta t_2}\,db_2 + \int_{\mathbb{R}\setminus[-1,1]} \widehat{\rho}(s_1,-\delta\frac{1}{|b_2|^\beta})e^{ib_2t_2}e^{-\delta t_2/|b_2|^\beta}\,db_2 \right) \end{aligned}$$

is, as a function of s_1, an analytic function on $U \times V$ and satisfies

$$\begin{aligned}|\widetilde{\rho}(s_1, t_2)| &\leq C\frac{\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}}{(1+|b_1|)^{M-D_0}}\left(2e^{-\delta t} + \int_{\mathbb{R}\setminus[-1,1]} \frac{e^{-\delta t/|b|^{\beta}}}{(1+|b|)^{M-D_0}}\, db\right) \\ &\leq C\frac{\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}}{(1+|b_1|)^{M-D_0}}(1+t_2)^{-\beta},\end{aligned}$$

as soon as $M > D_0 + 2$. We are interested in $\rho(t_1, t_2) = \frac{1}{4\pi^2}\int_{\mathbb{R}} \widetilde{\rho}(s_1, t_2)e^{-ib_1t_1}\, db_1$. In the same way, by moving the curve of integration in s_1 to Γ, we obtain (recall that we have assumed that $\int f\, dm_\lambda = \int g\, dm_\lambda = 0$):

$$\rho(t_1, t_2) \leq C\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}[(1+t_1)^{\beta} + (1+t_2)^{\beta}].$$

Observe that the above proof also yields, setting $u = 1$ for instance:

Proposition 7.13. *Let $\varphi_0 \in \mathcal{K}_\alpha(\Sigma_+)$as above. For $\varepsilon, C, c > 0$ and M as above, for all $\varphi, \|\varphi - \varphi_0\| < \varepsilon$, all $f, h \in \mathcal{K}^+_{\alpha,M}$, we have, for all positive t:*

$$\left|\rho_{f,h,m_\lambda}(t)\right| \leq C\|f\|_{\alpha,M}\|h\|_{\alpha,M}[(1+t)^{-c}]. \tag{7.10}$$

Indeed, if we assume $\int f\, dm_\varphi = 0$, this is exactly the same computation, with only one variable s. But (7.10) holds for f as soon as it holds for $f - \int f\, dm_\varphi$. By the same token, using Proposition 7.13, we can replace in (7.8) f and h by $f - \int f\, dm_\varphi$ and $h - \int h\, dm_\varphi$. This achieves the proof of Proposition 7.11. □

7.2.3. *From one-sided to two-sided smooth functions.* This part goes back to Ruelle ([**R**]), we present it here for completeness. We consider a new space of functions: for $\alpha > 0$ and $M \in \mathbb{N}$, let $\mathcal{K}'_{\alpha,M}$ be the set of functions f on Σ^τ with the following properties:

- for all $x \in \Sigma$, $f(x, r) = 0$ for r outside the interval $[\frac{\inf \tau}{3}, \frac{2\inf\tau}{3}]$,
- for all $x \in \Sigma$, $r \mapsto f(x, r)$ is of class C^M and
- the functions $\frac{\partial^k f}{\partial r^k}(x, r)$, for $0 \leq k \leq M$ are α-Hölder continuous on Σ and continuous in r.

For $f \in \mathcal{K}'_{\alpha,M}$, we still denote $\|f\|_{\alpha,M} := \sup_{r,k\leq M}\|\frac{\partial^k f}{\partial r^k}(., r)\|_\alpha$. We show in this subsection

Proposition 7.14. *There exist $\varepsilon', C', c' > 0$ and M such that for all φ such that $\|\varphi - \varphi_0\|_\alpha < \varepsilon'$, all $f, u, h \in \mathcal{K}'_{\alpha,M}$, we have, for all positive t_1, t_2:*

$$\left|\rho_{f,u,h,m_\varphi}(t_1, t_2)\right| \leq C'\|f\|_{\alpha,M}\|u\|_{\alpha,M}\|h\|_{\alpha,M}[(1+t_1)^{-c'} + (1+t_2)^{-c'}].$$

Proof. Assume first that $\int f\, dm_\lambda = \int u\, dm_\lambda = \int h\, dm_\lambda = 0$.

The following construction reduces the proof of Proposition 7.14 to a direct extension of the proof of Proposition 7.11. Let $A(x)$ be a function in $\mathcal{K}'_\alpha(\Sigma)$; then (see e.g. [**P1**]), there exists a decomposition $A = \sum_{j=0}^{\infty} A_j$, where

(1) $x \mapsto A_j(x)$ depends only on the coordinates $(x_{-j}, x_{-j+1}, \cdots)$ of x,
(2) $\sup_x |A_j(x)| \le \alpha^j \|A\|_\alpha$ and
(3) $\|A_j\|_\alpha \le \|A\|_\alpha$.

We also associate to A and $s \in \mathbb{C}$ the functions $\widetilde{A}_j^s(x) := e^{-s\tau^j(x)} A_j(\sigma^j x)$. Then, by [**R**] (see also [**D1**] and [**P1**]), there is $\alpha', 0 < \alpha' < \alpha$, and $\theta, 0 < \theta < 1$, such that, with $s = a + ib$,

(1) $x \mapsto \widetilde{A}_j^s(x)$ depends only on the coordinates $(x_0, x_1, \cdots)$ of x,
(2) $\sup_x |\widetilde{A}_j^s(x)| \le e^{Cj|a|} \alpha^j \|A\|_\alpha$ and
(3) $\|\widetilde{A}_j^s\|_{\alpha'} \le C e^{Cj|a|} |b| \theta^j \|A\|_\alpha$.

Finally, we set $\widetilde{A}^s(x) := \sum_j \widetilde{A}_j^s(x)$; we have, if $|a|$ is small enough,

(1) $x \mapsto \widetilde{A}^s(x)$ depends only on the coordinates $(x_0, x_1, \cdots)$ of x,
(2) $\sup_x |\widetilde{A}^s(x)| \le C \|A\|_\alpha$,
(3) $\|\widetilde{A}^s\|_{\alpha'} \le C|b| \|A\|_\alpha$ and
(4) $\int \widetilde{A}^0(x)\, d\overline{\nu}(x) = \int A(x)\, d\overline{\nu}(x)$ for any shift invariant measure $\overline{\nu}$ on Σ.

Property (4) is clear since $\widetilde{A}^0(x) = \sum_j \widetilde{A}_j^0(x) = \sum_j A_j(\sigma^j x)$, whereas $A(x) = \sum_j A_j(x)$ and both series of functions converge uniformly.

Choose ε' so that for all $\varphi, \|\varphi - \varphi_0\|_\alpha < \varepsilon'$. Proposition 7.6 and Proposition 7.10 apply on $\mathcal{K}_{\alpha'}$. Fix $f, u, h, \varphi \in \mathcal{K}_{\alpha,M}$ and write $\rho(t_1, t_2)$ for $\rho_{f,h,g,m_\varphi}(t_1, t_2)$ We now write as before the Laplace transform $\widehat{\rho}(s_1, s_2)$ of $\rho(t_1, t_2)$ as:

$$\widehat{\rho}(s_1, s_2) = \sum_{n,m} \int_\Sigma H(\sigma^{n+m} x, -s_2) e^{s_2 \tau^{m+n}(x)} U(\sigma^n x, s_2 - s_1) e^{(s_1 - s_2)\tau^n(x)} F(x, s_1)\, d\overline{\nu}_\phi(x).$$

We consider it as a series in the sense of tempered distributions: for any $B(s,t)$ in the Schwartz space of $\mathbb{R}^2$, $\int \widehat{B}(ib_1, ib_2) \widehat{\rho}(ib_1, ib_2)\, db_1 db_2$ makes sense and is equal to $-4\pi^2 \int B(t_1, t_2) \rho(t_1, t_2)\, dt_1 dt_2$. The series of integrals $\int B(t_1, t_2) \rho_{n,m}(t_1, t_2)\, dt_1 dt_2$ converges absolutely. It still does if one consider the sum over n, m in $\mathbb{Z}$ instead of $\mathbb{Z}_+$. For each $(n, m) \in \mathbb{Z} \times \mathbb{Z}$, we write, using the decompositions $H(x, s) = \sum_j H_j(x, s)$,

$U(x,s) = \sum_k U_k(x,s), F(x,s) = \sum_\ell F_\ell(x,s)$ and the above $\widetilde{A}_j^s$ notation:

$$\begin{aligned}
&\widehat{\rho}_{n,m}(s_1,s_2) = \\
= \ & \int_\Sigma H(\sigma^{n+m}x, -s_2)e^{s_2\tau^{m+n}(x)}U(\sigma^n x, s_2-s_1)e^{(s_1-s_2)\tau^n(x)}F(x,s_1)\, d\overline{\nu}_\phi(x) \\
= \ & \sum_{j,k,\ell}\int_\Sigma H_j(\sigma^{n+m}x, -s_2)e^{s_2\tau^{m+n}(x)}U_k(\sigma^n x, s_2-s_1)e^{(s_1-s_2)\tau^n(x)}F_\ell(x,s_1)\, d\overline{\nu}_\phi(x) \\
= \ & \sum_{j,k,\ell}\int_\Sigma \widetilde{H}_j^{-s_2}(\sigma^{n+m-j}x, -s_2)e^{s_2\tau^{m+n-j}(x)}\widetilde{U}_k^{(s_2-s_1)}(\sigma^{n-k}x, s_2-s_1)e^{(s_1-s_2)\tau^{n-k}(x)} \\
& \qquad\qquad \widetilde{F}_\ell^{s_1}(\sigma^{-\ell}x, s_1)e^{s_1\tau^\ell(\sigma^{-\ell}x)}\, d\overline{\nu}_\phi(x) \\
= \ & \sum_{j,k,\ell}\int_\Sigma \widetilde{H}_j(\sigma^{n+m-j}x, -s_2)e^{s_2\tau^{m+k-j}(\sigma^{n-k}x)}\widetilde{U}_k(\sigma^{n-k}x, s_2-s_1)e^{s_1\tau^{n-k+\ell}(\sigma^{-\ell}x)} \\
& \qquad\qquad \widetilde{F}_\ell(\sigma^{-\ell}x, s_1)\, d\overline{\nu}_\phi(x),
\end{aligned}$$

where we used the cocycle relation $\tau^{n+m}(x) = \tau^n(x) + \tau^m(\sigma^n x)$ valid for all $m,n \in \mathbb{Z}$ and wrote $\widetilde{F}(x,s)$ for $\widetilde{F}^s(x,s)$, $\widetilde{G}_j(x,s)$ for $\widetilde{G}_j^s(x,s)$ and $\widetilde{U}_k(x,s)$ for $\widetilde{U}_k^s(x,s)$.

We now replace the summation in (n,m) by a summation in (p,q), where $p := n-k+\ell, q := m+k-j$. Assume for example $p \geq 0, q \geq 0$ (and then $p+q = n+m-j+l \geq 0$). We write, using the invariance of $\overline{\nu}_\phi$, the integral

$$\int_\Sigma \widetilde{H}_j(\sigma^{n+m-j}x, -s_2)e^{s_2\tau^{m+k-j}(\sigma^{n-k}x)}\widetilde{U}_k(\sigma^{n-k}x, s_2-s_1)e^{s_1\tau^{n-k+\ell}(\sigma^{-\ell}x)}\widetilde{F}_\ell(\sigma^{-\ell}x, s_1)\, d\overline{\nu}_\phi(x), \tag{7.11}$$

as:

$$\int_\Sigma \widetilde{H}_j(\sigma^{n+m-j+\ell}x, -s_2)e^{s_2\tau^{m+k-j}(\sigma^{n-k+\ell}x)}\widetilde{U}_k(\sigma^{n-k+\ell}x, s_2-s_1)e^{s_1\tau^{n-k+\ell}(x)}\widetilde{F}_\ell(x, s_1)\, d\nu_\phi(x),,$$

where we replaced $\overline{\nu}_\phi$ by ν_ϕ since the integrand now depends only on the non-negative coordinates of x. As before, we can write these integrals using the transfer operators as

$$\begin{aligned}
& \textstyle\int_\Sigma \widetilde{H}_j(\sigma^{m+k-j}x, -s_2)e^{s_2\tau^{m+k-j}(x)}\widetilde{U}_k(x, s_2-s_1)\mathcal{L}_{\phi+s_1\tau}^{n-k+\ell}(\widetilde{F}_\ell(.,s_1))(x)\, d\nu_\lambda(x) \\
= \quad & \textstyle\int_\Sigma \widetilde{H}_j(x, -s_2)\mathcal{L}_{\phi+s_2\tau}^q[\widetilde{U}_k(.,s_2-s_1)\mathcal{L}_{\phi+s_1\tau}^p(\widetilde{F}_\ell(..,s_1))(.)](x)\, d\nu_\lambda(x).
\end{aligned}$$

If $|a_1|, |a_2|$, and $|a_1-a_2|$ are small enough, one can sum in $j,k,\ell \in \mathbb{Z}_+^3$ the integral (7.11) for the same value of (p,q);we obtain, when $p,q \geq 0$,

$$\int_\Sigma \widetilde{H}(x, -s_2)\mathcal{L}_{\phi+s_2\tau}^q[\widetilde{U}(.,s_2-s_1)\mathcal{L}_{\phi+s_1\tau}^p(\widetilde{F}(..,s_1))(.)](x)\, d\nu_\phi(x).$$

The other possible signs of p,q and $p+q$ are treated in the same way.

By applying Proposition 7.6 to $\mathcal{K}_{\alpha'}$, we conclude that there are positive numbers δ', β' such that, for all $\varphi, \|\varphi-\varphi_0\|_{\alpha'} < \varepsilon'$ the series of operators $\sum_n \mathcal{L}_{\phi+s\tau}^n$ converges and is analytic on the region U'

$$U' \ := \ \{s = a+ib; |b| > 1 \ |a| < \frac{2\delta'}{|b|^{\beta'/2}}\}$$

and that there is a constant D_0' such that, for $s \in U'$,

$$\| \sum_n \mathcal{L}^n_{\phi+s\tau} \| \leq D_0' |b|^{D_0'}. \tag{7.12}$$

Moreover, there is $\delta' > 0$ such that on the series of operators $\sum_n \mathcal{L}^n_{\phi+s\tau}$ converges and is meromorphic on the region V',

$$V' := \{s, s = a + ib; |b| < 2, \ |a| < 2\delta'\}$$

with a simple pole at 0 and residue the projection on the constant function $\nu_\phi(.)$. We conclude as above (but with a different argument for each one of the six sums over $(p,q), (-q, p+q), (-p, p+q), (-p-q, q), (p, -p-q), (-p, -q)$ in $(\mathbb{Z}_+ \times \mathbb{Z}_+)$) that $\widehat{\rho}(s_1, s_2)$ is given by an analytic function defined on the region where s_1, s_2 and $s_1 - s_2$ all belong to $U' \cup V'$ (and have a real part smaller than δ_0) and satisfying

$$|\widehat{\rho}(s_1, s_2)| \leq C\|h\|_{\alpha,M}\|u\|_{\alpha,M}\|f\|_{\alpha,M}(1+|b_1|)^{D_0''-M}(1+|b_1-b_2|)^{D_0''-M}(1+|b_2|)^{D_0''-M},$$

where $D_0'' = D_0' + 1$.

If M has been chosen greater than $D_0'' + 2$, we obtain Proposition 7.14 (for functions with integral 0) by the same argument as before, provided one chooses in each of the six cases contours Γ of integration with the right sign.

The extension of Proposition 7.13 to functions $f, h \in \mathcal{K}'_{\alpha,M}$ with $\int f dm_\phi = 0, \int h dm_\phi = 0$ goes again by the same computation, without the function u. Again, (7.10) holds for f as soon as it holds for $f - \int f \, dm_\varphi$. This justifies the reduction to functions with integral 0 in the proof of proposition 7.14. $\square$

7.2.4. *Hölder continuous functions.* We conclude the proof of Theorem 7.2 and of Proposition 7.3 by approximating any Hölder continuous function by regular functions. We have proven (7.1) for functions in $\mathcal{K}'_{\alpha,M}$ with some constants C', c'; (7.1) holds also if f, u, h are such that $f \circ \sigma_{t_1}, u \circ \sigma_{t_2}, h \circ \sigma_{t_3} \in \mathcal{K}'_{\alpha,M}$ for bounded $t_i, i = 1,2,3$. There is $C_9 = 10 + 6\frac{\sup_x \tau(x)}{\inf_x \tau(x)}$ such that any function which is of class C^M along the trajectories of the special flow $(\Sigma^\tau, \sigma_t, t \in \mathbb{R})$ and such that the first M derivatives along the flow are α-Hölder continuous functions can be written as a sum of less than C_9 functions in $\mathcal{K}'_{\alpha,M}$. Using the projection from the manifold to Σ^τ, we conclude that there exist $\varepsilon, C'', c' > 0, \alpha, M$ such that for all for all $\varphi, \|\varphi - \varphi_0\|_\alpha < \varepsilon$, all f, u, h that are of class C^M along the trajectories of the flow and such that all the derivatives along the flow up to order M belongs to $\mathcal{K}_\alpha(SM)$, we have, for all $t_1, t_2 \geq 0$:

$$\left|\rho_{f,u,h,m_\varphi}(t_1, t_2)\right| \leq C''\|f\|_{\alpha,M}\|u\|_{\alpha,M}\|h\|_{\alpha,M}[(1+t_1)^{-c'} + (1+t_2)^{-c'}],$$

where $\|.\|_{\alpha,M}$ is the maximum of the $\|\|_\alpha$ norms of the first M derivatives along the flow.

We conclude by smoothing all functions in $\mathcal{K}_\alpha$. Let $\overline{\psi}$ be a C^M nonnegative function on $\mathbb{R}$, with support in $[-1, +1]$ and integral 1. For $\varepsilon > 0$ and a function $f \in \mathcal{K}_\alpha$, set

$$\overline{\psi}_\varepsilon(t) := \frac{1}{\varepsilon}\overline{\psi}(\frac{t}{\varepsilon}) \text{ and } f_\varepsilon(x) := \int_{\mathbb{R}} \overline{\psi}_\varepsilon(t) f(\varphi_t x) \, dt.$$

We have $\sup_x |f(x) - f_\varepsilon(x)| \le \varepsilon^\alpha \|f\|_\alpha$ and $\|f_\varepsilon\|_{\alpha,M} \le \varepsilon^{-M-1}\|f\|_\alpha$.

Fix $t_1, t_2 > 0$, choose $\varepsilon = [1/3(1+t_1)^{-c'} + 1/3(1+{}_2)^{-c'}]^{\frac{1}{\alpha+3M+3}}$ and replace f, u, h by $f_\varepsilon, u_\varepsilon, h_\varepsilon$. One obtains (7.1) for f, u, h with some constant C_0' and $c_0' = \frac{c'\alpha}{\alpha+3M+3}$.

8. Appendix 2

In this section, we prove the well-known decomposition results of the Green function on a barrier in the form that we use in Section 3. Let $\widetilde{M}$ be a simply connected nonpositively curved Hadamard manifold with bounded geometry. We denote λ_0 the bottom of the spectrum of the Laplacian and we assume that the Green function G_{λ_0} is finite. Let $\mathcal{D}$ be an open domain in $\widetilde{M}$. Recall that for $\lambda \le \lambda_0$ the relative Green function $G_\lambda(x, y : \mathcal{D})$ defined as the Green function of the operator $\Delta + \lambda$ on $\mathcal{D}$ with Dirichlet conditions at the boundary.

Consider the Brownian motion on $\widetilde{M}$ and for $x \in \widetilde{M}$, denote by $\mathbb{P}_x$ the probability measure it defines on the set Ω_x of continuous paths starting at x, $\mathbb{E}_x$ the associated expectation. Let A be a closed subset of $\widetilde{M}$ and assume $x \notin A$. For $\omega \in \Omega_x$, let $T_A(\omega) \in]0, +\infty]$ be the first time the trajectory ω hits A. In probabilistic terms, we have for every nonnegative measurable function F,

$$\int_{\widetilde{M}\setminus A} F(y) G_\lambda(x, y : \widetilde{M} \setminus A)dy \;=\; \mathbb{E}_x \left[\int_0^{T_A(\omega)} e^{\lambda t} F(w_t)\, dt \right]. \tag{8.1}$$

Consider A an $(n-1)$-dimensional submanifold, with bounded geometry and assume $x, z \in \mathcal{D}$. Write $T(\omega)$ for $T_{A\cup\partial\mathcal{D}}(\omega)$. Observe that if $T(\omega) < T_{\partial\mathcal{D}}(\omega)$, $\omega_{T(\omega)} \in A \cap \mathcal{D}$. In particular, in that case, $G_\lambda(\omega_{T(\omega)}, z : \mathcal{D})$ makes sense. We have, for all $\lambda \le \lambda_0$,

Proposition 8.1.

$$G_\lambda(x, z : \mathcal{D}) \;=\; \mathbb{E}_x \left[1_{T(\omega)<T_{\partial\mathcal{D}}(\omega)} e^{\lambda T(\omega)} G_\lambda(\omega_{T(\omega)}, z : \mathcal{D}) \right] + G_\lambda(x, z : \mathcal{D} \setminus A).$$

Proof. Indeed, if $\delta < d(z, A \cup \partial\mathcal{D})$, we may write

$$\begin{aligned}
&\int_{B(z,\delta)} G_\lambda(x, w : \mathcal{D})\, dw \\
=\;& \mathbb{E}_x \left[\int_0^{T_{\partial\mathcal{D}}(\omega)} e^{\lambda t} 1_{B(z,\delta)}(\omega_t)\, dt \right] \\
=\;& \mathbb{E}_x \left[1_{T(w)<T_{\mathcal{D}}(w)} \int_{T(w)}^{T_{\mathcal{D}}(w)} e^{\lambda t} 1_{B(z,\delta)}(w_t)\, dt \right] + \mathbb{E}_x \left[1_{T(w)<T_{\mathcal{D}}(w)} \int_0^{T(w)} e^{\lambda t} 1_{B(z,\delta)}(w_t)\, dt \right] \\
&+\mathbb{E}_x \left[1_{T(w)\ge T_{\mathcal{D}}(w)} \int_0^{T_{\mathcal{D}}(w)} e^{\lambda t} 1_{B(z,\delta)}(w_t)\, dt \right] \\
=\;& \mathbb{E}_x \left[1_{T(\omega)<T_{\partial\mathcal{D}}(\omega)} e^{\lambda T(\omega)} \int_{B(z,\delta)} G_\lambda(\omega_{T(\omega)}, w : \mathcal{D})\, dw \right] + \int_{B(z,\delta)} G_\lambda(x, y : \mathcal{D} \setminus A)dy.
\end{aligned}$$

We used the Strong Markov Property of the stopping time $T(\omega)$ to write the last line. The proposition follows by letting $\delta \to 0$. □

Let ϖ_x^λ be the distribution on $A \cap \mathcal{D}$ such that the proposition writes, for all $\lambda \leq \lambda_0$,

$$G_\lambda(x, z : \mathcal{D}) \;=\; \int_{A \cap \mathcal{D}} G_\lambda(y, z : \mathcal{D})\, d\varpi_x^\lambda(y) + G_\lambda(x, z : \mathcal{D} \setminus A)$$

(ϖ_x^0 is the distribution of the hitting point $\omega_{T(\omega)}$ on $A \cap \mathcal{D}$). The measure ϖ_x^λ has a density with respect to the induced Lebesgue measure dy on A. We now prove the estimate in Propositions 3.2 and 3.3 that we recall:

Proposition 8.2. *Let A be a $(n-1)$ dimensional submanifold of $\mathcal{D}$ with bounded geometry. Set A_1 for the set of points of A at distance at least 1 from $\mathcal{D}^c$. There exists a constant C_3 such that, for $\lambda \in [0, \lambda_0]$, for any positive function F on A_1 and any $x \in \mathcal{D}, d(x, A) > 1$,*

$$\int_{A_1} F(y) d\varpi_x^\lambda(y) \leq \; C_3 L(F)^2 \int_A G_\lambda(x, y) F(y)\, dy,$$

where $L(F) := e^{\sup_A ||\nabla \ln F||}$ is the (multiplicative) Lipschitz constant of F.

Proof. We choose a cover of A_1 by open balls $B(y_p, \delta), y_p \in A_1$ with bounded multiplicity C_5 (in $\widetilde{M}$) and a partition of unity φ_p on A_1 subordinate to the cover $B(y_p, \delta) \cap A_1$ of A_1 for some $\delta, 0 < \delta \leq 1/2$. We have to estimate:

$$\int_{A_1} F(y) d\varpi_x^\lambda(y) \leq \; \sum_k \sum_p e^{(k+1)\lambda} \mathbb{E}_x \left[1_{T(\omega) \in [k, k+1)} 1_{T(\omega) < T_{\mathcal{D}}(\omega)} \varphi_p(\omega_{T(\omega)}) F(\omega_{T(\omega)}) \right].$$

Firstly, we estimate from above F on $B(y_p, \delta)$ by $L(F) F(y_p)$.

Then, we write for all $s, k+2 \leq s < k+3$,

$$\begin{aligned} & \mathbb{P}_x \left[\omega_s \in B(y_p, \delta) \cap \mathcal{D}, s < T_{\mathcal{D}}(\omega) \right] \\ \geq \;& \mathbb{P}_x \left[\omega_s \in B(y_p, \delta) \cap \mathcal{D}, k \leq T(\omega) < k+1, s < T_{\mathcal{D}}(\omega), \omega_{T(\omega)} \in B(y_p, \delta) \cap A_1 \cap \mathcal{D} \right] \\ \geq \;& \mathbb{E}_x \left[1_{[k, k+1)}(T(\omega)) 1_{T(\omega) < T_{\mathcal{D}}(\omega)} 1_{B(y_p, \delta) \cap A_1 \cap \mathcal{D}}(\omega_{T(\omega)}) U(y_p, \omega_{T(\omega)}, s - T(\omega)) \right], \end{aligned}$$

where

$$U(y, z, t) \;:=\; \mathbb{P}_z \left[\omega_t \in B(y, \delta) \cap \mathcal{D}, 1 \leq t \leq T_{\mathcal{D}}(\omega) \right].$$

Here, we used the Strong Markov property to write the second inequality. Set

$$C_{10}^{-1} \;:=\; \inf\{U(y, z, t); y, z \in \mathcal{D}, d(y, z) \leq \delta, d(y, \mathcal{D}) > 1, 1 \leq t \leq 3\}.$$

The constant C_{10} is finite by bounded geometry and we have

$$\begin{aligned} & \mathbb{P}_x \left[\omega_s \in B(y_p, \delta) \cap \mathcal{D}, s < T_{\mathcal{D}}(\omega) \right] \\ \geq \;& C_{10}^{-1} \mathbb{P}_x \left[k \leq T(\omega) < k+1, T(\omega) < T_{\mathcal{D}}(\omega), \omega_{T(\omega)} \in B(y_p, \delta) \cap A_1 \cap \mathcal{D} \right]. \end{aligned}$$

It follows that

$$e^{(k+1)\lambda} \mathbb{E}_x \left[1_{T(\omega) \in [k, k+1)} 1_{T(\omega) < T_{\mathcal{D}}(\omega)} \varphi_p(\omega_{T(\omega)}) \right] \; \leq C_{10} \int_{k+2}^{k+3} \mathbb{E}_x \left[e^{\lambda s} 1_{B(y_p, \delta) \cap \mathcal{D}}(\omega_s) \right] ds.$$

We thus have, by summing over $k \in \mathbb{N}$,

$$\begin{aligned}
&\int_{A_1} F(y) d\varpi_x^\lambda(y) \\
\leq\ & C_{10} L(F) \sum_p F(y_p) \mathbb{E}_x \left[\int_0^\infty e^{\lambda s} 1_{B(y_p,\delta)\cap\mathcal{D}}(\omega_s)\, ds \right] \\
\leq\ & C_{10} L(F) \sum_p F(y_p) \int_{B(y_p,\delta)\cap\mathcal{D}} G_\lambda(x,w)\, dw.
\end{aligned}$$

By bounded geometry, we can choose δ small enough that for $d(w, A_1) < \delta$, the orthogonal projection of w on A is well defined. By Harnack inequality and the Lipschitz regularity of F, we have:

$$\begin{aligned}
\int_{A_1} F(y) d\varpi_x^\lambda(y) &\leq C_{10} C_0 L(F)^2 \sum_p \int_{B(y_p,\delta)} F(\overline{w}) G_\lambda(x,\overline{w})\, dw \\
&\leq C_5 C_{10} C_0 L(F)^2 \int_{\cup_p B(y_p,\delta)} F(\overline{w}) G_\lambda(x,\overline{w})\, dw.
\end{aligned}$$

The inequality follows, with one more geometric constant C_{11} such that for any continuous nonnegative function G on $\partial\mathcal{C}$, $\int_{\cup_{y\in A_1} B(y,\delta)} G(\overline{w}) dw \leq C_{11} \int_A G(y) dy$. ☐

Proposition 8.3. *Let $\mathcal{C}$ be an open domain, $\mathcal{C} \subset \mathcal{D}, d(\mathcal{C}, \partial\mathcal{D}) > 1$. Let $x \in \mathcal{C}$, $A := \partial\mathcal{C}$ and let ϖ_x^λ be the distribution in Proposition 3.1 on A. Then, there exists a constant C_3 such that if $d(x, A) > 1$, then for all $\lambda \in [0, \lambda_0]$,*

$$C_3 (L(F))^{-2} \int_A G_\lambda(x, y : \mathcal{D}) F(y)\, dy \ \leq\ \int_A F(y) d\varpi_x^\lambda(y).$$

Proof. The proof is similar to the proof of Proposition 3.2. We write:

$$\begin{aligned}
&\mathbb{E}_x \left[\int_0^{T_{\mathcal{D}}(\omega)} e^{\lambda s} 1_{B(\omega_{T(\omega)},3/4)\cap(\mathcal{D}\setminus\mathcal{C})}(\omega_s) F(\omega_{T(\omega)})\, ds \right] \\
=\ & \mathbb{E}_x \left[e^{\lambda T(\omega)} F(\omega_{T(\omega)}) \mathbb{E}_{\omega_{T(\omega)}} \left[\int_0^{T_{\mathcal{D}}(\omega_{T(\omega)})} e^{\lambda(s-T(\omega)} 1_{B(\omega_{T(\omega)},3/4)\cap(\mathcal{D}\setminus\mathcal{C})}(\omega_{s-T(\omega)})\, ds \right] \right] \\
\leq\ & C_3^{-1} \mathbb{E}_x \left[e^{\lambda T(\omega)} F(\omega_{T(\omega)}) \right],
\end{aligned}$$

where $c_{10}^{-1} = \sup_{y \in A} \int_{B(y,3/4)\cap(\mathcal{D}\setminus\mathcal{C})} G_{\lambda_0}(y, z : \mathcal{D})\, dz$, which is finite by bounded geometry since $d(A, \mathcal{D}) > 1$.

It follows that

$$\begin{aligned}
\int_A F(y)d\varpi_x^\lambda(y) &= \mathbb{E}_x\left[e^{\lambda T(\omega)}F(\omega_{T(\omega)}1_{T(\omega)<\infty}\right] \\
\geq & \ c_{10}(L(F))^{-1}\sum_p F(y_p)\mathbb{E}_x\left[\int_0^{T_{\mathcal{D}}(\omega)} e^{\lambda s}1_{B(\omega_{T(\omega)},3/4)\cap(\mathcal{D}\setminus\mathcal{C})}(\omega_s)\varphi_p(\omega_{T(\omega)})\,ds\right] \\
\geq & \ c_{10}(L(F))^{-1}C_0^{-1}\sum_p F(y_p)\int_{B(y_p,3/4)\cap\mathcal{D}\setminus\mathcal{C}} G_\lambda(x,z:\mathcal{D})dz \\
\geq & \ C_0^{-1}L(F)^{-2}C_0^{-1}\int_{\{w:d(w,A)\leq 1/4\}} F(\overline{w})G_\lambda(x,\overline{w}:\mathcal{D})\,dw \\
\geq & \ C_0^{-1}L(F)^{-2}C_0^{-1}c_{13}\int_A F(y)G_\lambda(x,y:\mathcal{D})\,dy,
\end{aligned}$$

where c_{13} is another geometric constant. □

Observe that the geometry of A plays a role only in the choice of δ and of C_{11}, c_{13}. In particular, the estimates of Propositions 3.2 and 3.3 are uniform for all the barriers in the text.

REFERENCES

[**ABJ**] J.-P. Anker, P. Bougerol and T. Jeulin, *The infinite Brownian loop on a symmetric space,* Rev. Mat. Iberoam., **18** (2002), 41–97.

[**An1**] A. Ancona, *Negatively curved manifolds, elliptic operators and the Martin boundary*, Ann. Math. (2) **125** (1987) 495–536.

[**An2**] A. Ancona, *Théorie du potentiel sur les graphes et les variétés* in École d'Été de Probabilités de Saint-Flour XVIII 1988- edited by Paul-Louis Hennequin, Springer Lecture Notes, Berlin, Heidelberg, Springer-Verlag, **1427** (1990), 5–112.

[**Ano**] D. Anosov, *Theory of dynamical systems. Part I: Ergodic theory.*, Lectures held in Warsaw, Spring, 1967. Lecture Notes Series, No. 23 Matematisk Institut, Aarhus Universitet, Aarhus 1970.

[**AS**] D. Anosov, J. Sinai, *Certain smooth ergodic systems. (Russian)*, Uspehi Mat. Nauk **22** (1967) no. 5 (137), 107–172.

[**AnS**] M.T. Anderson and R. Schoen, *Positive harmonic functions on complete manifolds of negative curvature*, Annals Math. (2) **121** (1985), 429–461.

[**BL**] W. Ballmann and F. Ledrappier, *Discretization of positive harmonic functions on Riemannian manifolds and Martin Boundary* , in Actes de la table ronde de géométrie différentielle en l'honneur de Marcel Berger, Arthur L. Besse ed., Séminaires et Congrès, **1** (1996), 77-92.

[**B**] P. Bougerol, *Théorème central limite local sur certains groupes de Lie*, Ann. Sci. École Norm. Sup. (4) **14** (1981), 403–432.

[**Br**] R. Brooks, *The fundamental group and the spectrum of the Laplacian*, Comment. Math. Helv. **56** (1981), 581–598.

[**BR**] R. Bowen and D. Ruelle, *The ergodic theory of Axiom A flows,* Invent. Math., **29** (1975), 181–202.

[**CY**] S. Y. Cheng and S. T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Commun. Pure Appl. Math. **28** (1975) 333–354.

[**Da**] E. B. Davies, *Non-Gaussian aspects of heat kernel behavior*, J. London Mat. Soc. (2) **55** (1997), 105–125.

[**D1**] D. Dolgopyat, *On decay of correlations in Anosov flows,* Ann. Math., **147** (1998), 357–390.

[**D2**] D. Dolgopyat, *Prevalence of rapid mixing in hyperbolic flows,* Ergod. Th. & Dynam. Sys. **18** (1998), 1097–1114.

[DGM] A. Debiard, B. Gaveau and E. Mazet, *Théorèmes de comparaison en géométrie riemannienne*, Publ. Kyoto Univ. **12** (1976), 391–425.

[F] W. Feller, *An introduction to probability theory and its applications. Vol. II. John Wiley & Sons.*

[Ge] P. Gerl, *Ein Gleichverteilungsatz auf F_2, in* Probability Measures on Groups, Springer Lecture Notes in Math. **706** Springer-Verlag, Berlin-Heidelberg-New York, (1979), 126–130.

[G1] S. Gouëzel, *Local limit theorem for symmetric random walks in Gromov-hyperbolic groups*, J. Amer. Math. Soc., **27** (2014), 893–928.

[G2] S. Gouëzel, *Martin boundary of random walks with unbounded jumps in hyperbolic groups*, Ann. Proba., **43** (2015), 2374–2404.

[GL] S. Gouëzel and S. Lalley, *Random walks on co-compact Fuchsian groups*, Ann. Sci. École Norm. Sup. (4) **46** (2013).

[GW] P. Gerl and W. Woess, *Local limits and Harmonic Functions for Nonisotropic Random Walks on Free Groups,* Proba. Th. Rel. Fields **71** (1986), 341–355.

[H1] U. Hamenstädt, *An explicite description of harmonic measure*, Math. Z. **205** (1990), 287–299.

[H2] U. Hamenstädt, *Harmonic measures for compact negatively curved manifolds*, Acta Math. **178** (1997), 39-107.

[H3] U. Hamenstädt, *Harmonic measures, Hausdorff measures and positive eigenfunctions*, J. Diff. Geom. **44** (1996), 1–31.

[Ka] T. Kato, Perturbation theory of linear operators, *Grund. math. Wissen.* **132** Springer Verlaf, Berlin-Heidelberg-New York (1980).

[K1] V. A. Kaimanovich, *Invariant measures of the geodesic flow and measures at infinity on negatively curved manifolds*, Ann. Inst. H. Poincaré, A, Phys. Théor., **53** (1990), 361–393.

[Ko] G. Kozma, *A graph counter-example to Davies's conjecture,* Rev. Mat. Iberoam. **30** (2014), 1–12.

[La] S. Lalley, *Finite range random walks on free groups and homogeneous trees*, Ann. Prob. **21** (1993), 2087–2130.

[L1] F. Ledrappier, *A renewal theorem for the distance in negative curvature*, Stochastic analysis (Ithaca, New York, 1993) Proc. Symp. Pure Math. **57** (1995), 351–360.

[L2] F. Ledrappier, *Structure au bord des variétés à courbure négative,* Séminaire Théorie Spec. Géom. Grenoble, **13** (1995), 97-122.

[L3] F. Ledrappier, *Applications of dynamics to compact manifolds of negative curvature*, *Proceedings of the International Congress of Mathematicians, Vol 1,2 (Zürich, 1994)*, Birkhäuser, Basel (1995), 1195–1202.

[Li] C. Liverani, *On Contact Anosov flows,* Annals of Math., **159** (2004), 1275–1312.

[M] J. Moser, *On Harnack theorem for elliptic differential equations*, *Comm. Pure Appl. Math.* **14** (1961), 577-591.

[M1] G. Margulis, *Applications of ergodic theory to the investigation of manifolds of negative curvature*, *Functional Anal. Appl.* **3** (1969), 335–336.

[M2] G. Margulis, On Some Aspects of the Theory of Anosov Systems, *Springer Monographs in Math.* Springer-Verlag, Berlin Heidelberg (2004).

[Me] I. Melbourne, *Rapid decay of correlations for nonuniformly hyperbolic flows,* Trans. AMS, **359**, (2007), 2421–2441.

[Mo] O. Mohsen, *Le bas du spectre d'une variété hyperbolique est un point selle*, Ann. Sci. École Norm. Sup. (4), **40** (2007), 191–207.

[Mv] S. A. Molchanov, *Diffusion processes and Riemannian geometry,* Russian Math. Surveys **30** (1975), 1–63.

[N] L. Naïm, *Sur le rôle de la frontière de R.S. Martin dans la théorie du potentiel*, Ann. Inst. Fourier, Grenoble **7** (1957), 183–281.

[Pi] R.G. Pinsky , Positive Harmonic Functions and Diffusion, Cambridge University Press, Cambridge, (1995).

[P1] M. Pollicott, *Multiple mixing for hyperbolic flows, preprint*

[P2] M. Pollicott, *Uniform Dolgopyat super-polynomial mixing, in preparation; personal communication.*

[PP] W. Parry and M. Pollicott, Zeta functions and the periodic orbit structure of hyperbolic dynamics. *Astérisque* **187–188** (1990).

[**PPS**] F. Paulin, M. Pollicott, B. Schapira, Equilibrium states in negative curvature, *Astérisque*, **373** (2015).
[**R**] D. Ruelle, *Resonances for Axiom A flows,* J. Diff. Geometry **25** (1987), 99–116.
[**SZ**] P. Souplet, Q. Zhang, *Sharp gradient estimates and Yau Liouville theorem for the heat equation on non-compact manifolds*, Bull. London Math. Soc. 38 (2006) 1045–1053.
[**S**] D. Sullivan, *Related aspects of positivity in Riemannian geometry,* J. Diff. Geom. **25** (1987), 327–351.
[**V**] N. Varopoulos, *Théorie du potentiel sur des groupes et des variétés*, C.R. Acad. Sci. Paris Sér. I Math. **302** (1986), 203–205.
[**W**] P. Walters, An introduction to ergodic theory, *Graduate texts in Mathematics* **79**, Springer-Verlag, New York-Berlin, (1982).

Department of Mathematics, 255 Hurley Hall, University of Notre Dame, Notre Dame IN, USA

Sorbonne Universités, UPMC Univ. Paris 06, UMR 7599, LPMA, F-75005, Paris, France

CNRS, UMR 7599, LPMA, F-75005, Paris, France

E-mail address: `ledrappier.1@nd.edu`

Department of Mathematical Sciences, Seoul National University, Seoul 151-747

E-mail address: `slim@snu.ac.kr`